

Rahul (Tony) Rao

Dual diagnosis: a call to action

Following publication of the last issue of *Advances in Dual Diagnosis*, we have launched a call for papers on topics that are often overlooked but have considerable relevance to dual diagnosis. In this editorial, we provide an overview of each topic so that they can serve as a springboard for the future. This is all the more relevant for a journal that spans the landscape of dual diagnosis across qualitative and empirical research, systematic and general reviews, viewpoint articles, conceptual papers and case studies. All our papers are required to show originality, preferably accompanied by research, practical and social implications. This strikes at the heart of an ethos that our papers have direct relevance to policy and practice. The topics covered in the call for papers cover intellectual disability (ID), novel psychoactive substances, pharmacological approaches and the interface with primary care.

Rahul (Tony) Rao is based at South London and Maudsley NHS Foundation Trust, London, UK, and Institute of Psychiatry Psychology and Neuroscience, London, UK.

Intellectual disability

People with ID have poorer health and social outcomes than those without, mainly through a lack of research to guide prevention and treatment. Up to 5% of people with ID have dual diagnosis ([van Duijvenbode et al., 2015](#)) but this group is also less likely to receive treatment of their dual diagnosis ([Chapman and Wu, 2012](#)). This may be an under-estimation as communication problems may increase the risk of poorer outcomes from pre-existing genetic disorders. This is compounded by the side effects from prescribed psychotropic medication and prescribed substances ([Degenhardt, 2000](#)). As such, dual diagnosis practitioners have a role in clinically effective service delivery ([Larson et al., 2017](#)). Public attitudes to the “double whammy” of ID and dual diagnosis can also reduce the likelihood of help-seeking through stigma ([Asamoah et al., 2009](#)), with subsequent under-reporting of substance use.

Forensic services represent a setting where people with ID are over-represented compared with non-forensic settings ([Raina and Lunsky, 2010](#)), with a clear association found between mild ID, offending and substance use ([Glaser and Florio, 2004](#)). There may be several factors underlying this association, including impaired judgement and risk taking from impulsivity ([Taylor, 2010](#)). Emotional lability mood swings with increased risk of self-harm are also known to be more likely in people with ID and dual diagnosis compared to those who do not use substances ([To et al., 2014](#)).

Novel psychoactive substances

There has been growing use of Novel Psychoactive Substances (NPS) over the past decade from their increasing availability through rogue websites and the “deep web”, with vulnerable groups such as younger people and those with a history of mental disorder at particular risk. Synthetic cannabinoids, cathinones, phenethylamines and dissociative substances such as ketamine account for the majority of NPS. These substances are often used in combination with others such as alcohol, cocaine and opioids. Higher levels of NPS use are known to exist in people with personality disorder, psychotic disorders and bipolar disorder. However, many NPS are undetected during urinary drug screening. More research into dual diagnosis with NPS use is needed to fill the gap in research and practice.

Pharmacological approaches

There is little literature on pharmacological treatment of dual diagnosis, with no consistent international guidelines to inform policy and practice (Vitali *et al.*, 2018). This is particularly stark in studies on best practice and treatment outcomes for people with psychotic disorders who have co-morbid alcohol use and the treatment of opioid use disorders in those with co-morbid depression (Murthy and Chand, 2012).

Interface with primary care

In primary care settings, dual diagnosis may be overlooked and masked by physical disorders. The detection of both substance use and mental disorders can help with both referral to drug and alcohol services and provide integrated rather than parallel or sequential care. This applies to both general (Ziedonis and Brady, 1997) and older populations (Rao *et al.*, 2019).

There is much food for thought in expanding the horizon of dual diagnosis and we hope that this call to action will not only improve research and practice in under-researched areas but improve the lives of people with dual diagnosis so that they receive the assessment, treatment and interventions they deserve in both developed and developing countries.

References

- Asamoah, G., Varughese, S., Mushtaq, S., Butterworth, L., Abraham, A. and Luty, J. (2009), "A randomised trial of ethnicity and stigmatised attitudes towards learning disability and alcoholism", *Ethnicity and Inequalities in Health and Social Care*, Vol. 2 No. 2, pp. 11-19, doi: [10.1108/17570980200900011](https://doi.org/10.1108/17570980200900011).
- Chapman, S.L.C. and Wu, L.T. (2012), "Substance abuse among individuals with intellectual disabilities", *Research in Developmental Disabilities*, Vol. 33 No. 4, pp. 1147-1156, doi: [10.1016/j.ridd.2012.02.009](https://doi.org/10.1016/j.ridd.2012.02.009).
- Degenhardt, L. (2000), "Interventions for people with alcohol use disorders and an intellectual disability: a review of the literature", *Journal of Intellectual & Developmental Disability*, Vol. 25 No. 2, pp. 135-146, doi: [10.1080/13269780050033553](https://doi.org/10.1080/13269780050033553).
- Glaser, W. and Florio, D. (2004), "Beyond specialist programmes: a study of the needs of offenders with intellectual disability requiring psychiatric attention", *Journal of Intellectual Disability Research*, Vol. 48 No. 6, pp. 591-602, doi: [10.1111/j.1365-2788.2004.00628.x](https://doi.org/10.1111/j.1365-2788.2004.00628.x).
- Larson, S., Eschenbacher, H., Anderson, L., Pettingell, S., Hewitt, A., Sowers, M., Bourne, M.L., Taylor, B. and Agosta, J. (2017), "In-home and residential long-term supports and services for persons with intellectual or developmental disabilities: status and trends through 2015", Residential Information Systems Project Report. Institute on Community Integration, available at: https://ici-s.umn.edu/files/aCHyYaFjMi/risp_2017
- Murthy, P. and Chand, P. (2012), "Treatment of dual diagnosis disorders", *Current Opinion in Psychiatry*, Vol. 25 No. 3, pp. 194-200, doi: [10.1097/YCO.0b013e328351a3e0](https://doi.org/10.1097/YCO.0b013e328351a3e0).
- Raina, P. and Lunsky, Y. (2010), "A comparison study of adults with intellectual disability and psychiatric disorder with and without forensic involvement", *Research in Developmental Disabilities*, Vol. 31 No. 1, pp. 218-223, doi: [10.1016/j.ridd.2009.09.008](https://doi.org/10.1016/j.ridd.2009.09.008).
- Rao, R., Crome, I., Crome, P. and Iliffe, S. (2019), "Substance misuse in later life: challenges for primary care: a review of policy and evidence", *Primary Health Care Research & Development*, Vol. 20 No. e117, pp. 1-7, doi: [10.1017/S1463423618000440](https://doi.org/10.1017/S1463423618000440).
- Taylor, J. (2010), "Psychotherapy for people with learning disabilities: creating possibilities and opportunities. A review of the literature", *Journal of Learning Disabilities and Offending Behaviour*, Vol. 1 No. 3, pp. 15-25, doi: [10.5042/jldob.2010.0625](https://doi.org/10.5042/jldob.2010.0625).
- To, W.T., Neiryck, S., Vanderplasschen, W., Vanheule, S. and Vandeveld, S. (2014), "Substance use and misuse in persons with intellectual disabilities (ID): results of a survey in ID and addiction services in Flanders", *Research in Developmental Disabilities*, Vol. 35 No. 1, pp. 1-9, doi: [10.1016/j.ridd.2013.10.015](https://doi.org/10.1016/j.ridd.2013.10.015).

van Duijvenbode, N., VanDerNagel, J.E., Didden, R., Engels, R.C., Buitelaar, J.K., Kiewik, M. and de Jong, C.A. (2015), "Substance use disorders in individuals with mild to borderline intellectual disability: current status and future directions", *Research in Developmental Disabilities*, Vol. 38 No. 3, pp. 319-328, doi: [10.1016/j.ridd.2014.12.029](https://doi.org/10.1016/j.ridd.2014.12.029).

Vitali, M., Mistretta, M., Alessandrini, G., Coriale, G., Romeo, M., Attilia, F., Rotondo, C., Sorbo, F., Pisciotta, F., Attilia, M.L. and Ceccanti, M. (2018), "Pharmacological treatment for dual diagnosis: a literature update and a proposal of intervention", *Rivista di Psichiatria*, Vol. 53 No. 3, pp. 160-169, doi: [10.1708/2925.29419](https://doi.org/10.1708/2925.29419).

Ziedonis, D. and Brady, K. (1997), "Dual diagnosis in primary care: detecting and treating both the addiction and mental illness", *Medical Clinics of North America*, Vol. 81 No. 4, pp. 1017-1036, doi: [10.1016/s0025-7125\(05\)70561-7](https://doi.org/10.1016/s0025-7125(05)70561-7).