

# The implicit midpoint rule for nonexpansive mappings in 2-uniformly convex hyperbolic spaces

H. Fukhar-ud-din and A.R. Khan

*Department of Mathematics and Statistics,  
King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia*

Implicit midpoint  
rule for  
nonexpansive  
mappings

95

Received 24 December 2018  
Revised 14 February 2019  
Accepted 17 February 2019

## Abstract

The purpose of this paper is to introduce the implicit midpoint rule (IMR) of nonexpansive mappings in 2-uniformly convex hyperbolic spaces and study its convergence. Strong and  $\Delta$ -convergence theorems based on this algorithm are proved in this new setting. The results obtained hold concurrently in uniformly convex Banach spaces, CAT(0) spaces and Hilbert spaces as special cases.

**Keywords** Uniformly convex hyperbolic space, Nonexpansive mapping, Midpoint rule, Fixed point, Condition(A), Convergence

**Paper type** Original Article

## 1. Introduction

The iterative methods for approximating fixed points of nonexpansive mappings have received a great attention due to the fact that in many practical problems, the controlling operators are nonexpansive (cf. [16]). The iterative methods of Mann [17] and Halpern [9] are very popular (see also [20]). An implicit iterative method was proposed [25] and studied in [7,12]. The IMR is a powerful numerical method for solving ordinary differential equations and differential algebraic equations. For related works in this context, we refer the reader to [2,5,20,22].

For the ordinary differential equation

$$y'(t) = g(t), \quad y_0 = y(0), \quad (1.1)$$

## JEL Classification — 47H09, 47H10

© H. Fukhar-ud-din and A.R. Khan. Published in the *Arab Journal of Mathematical Sciences*. Published by Emerald Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 4.0) license. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this license may be seen at <http://creativecommons.org/licenses/by/4.0/legalcode>

The authors are grateful to King Fahd University of Petroleum & Minerals (KFUPM) for supporting this research.

The publisher wishes to inform readers that the article “The implicit midpoint rule for nonexpansive mappings in 2-uniformly convex hyperbolic spaces” was originally published by the previous publisher of the *Arab Journal of Mathematical Sciences* and the pagination of this article has been subsequently changed. There has been no change to the content of the article. This change was necessary for the journal to transition from the previous publisher to the new one. The publisher sincerely apologises for any inconvenience caused. To access and cite this article, please use Fukhar-ud-din, H., Khan, A.R. (2019), “The implicit midpoint rule for nonexpansive mappings in 2-uniformly convex hyperbolic spaces” *Arab Journal of Mathematical Sciences*, Vol. 26 No. 1/2, pp. 95-105. The original publication date for this paper was 22/02/2019.



Arab Journal of Mathematical  
Sciences  
Vol. 26 No. 1/2, 2020  
pp. 95-105  
Emerald Publishing Limited  
e-ISSN: 2588-9214  
p-ISSN: 1319-5166  
DOI 10.1016/j.ajmsc.2019.02.002

IMR generates a sequence  $\{y_n\}$  via the relation

$$\frac{1}{h}(y_{n+1} - y_n) = g\left(\frac{y_{n+1} + y_n}{2}\right)$$

where  $h > 0$  is a step size. It is well known that if  $g: \mathbb{R}^k \rightarrow \mathbb{R}^k$  is Lipschitzian continuous and sufficiently smooth, then the sequence  $\{y_n\}$  converges to the exact solution of (1.1) as  $h \rightarrow 0$  uniformly over  $t \in [0, a]$  for any fixed  $a > 0$ .

Based on the above fact, Alghamdi et al. [1] presented the following IMR for nonexpansive mappings in the setting of a Hilbert space  $H$ :

$$y_{n+1} = (1 - t_n)y_n + t_n T\left(\frac{y_{n+1} + y_n}{2}\right) \tag{1.2}$$

where  $t_n \in (0, 1)$  and  $T: H \rightarrow H$  is a nonexpansive mapping and established weak convergence of (1.2) to the fixed point of  $T$  under some control conditions on  $\{t_n\}$ .

The extension of a linear version of a known result (usually in Banach spaces or Hilbert spaces) to metric spaces is very important. As an IMR for nonexpansive mappings involves general convex combinations, so we need some convex structure in a metric space to define an IMR on a nonlinear domain.

Let  $C$  be a nonempty subset of a metric space  $(M, d)$  and  $T: C \rightarrow C$  a mapping. Set  $F(T) = \{x \in M : Tx = x\}$ . The mapping  $T$  is: (i) nonexpansive if  $d(Tx, Ty) \leq d(x, y)$  for all  $x, y \in C$  (ii) quasi-nonexpansive if  $d(Tx, y) \leq d(x, y)$  for all  $x \in C$  and  $y \in F(T)$  (iii) semi-compact if for any bounded sequence  $\{x_n\}$  in  $C$  satisfying  $d(x_n, Tx_n) \rightarrow 0$ , there exists a subsequence  $\{x_{n_i}\}$  of  $\{x_n\}$  such that  $x_{n_i} \rightarrow x \in C$  (iv) completely continuous if every bounded sequence  $\{x_n\}$  in  $C$  implies that  $\{Tx_n\}$  has a convergent subsequence. A sequence  $\{x_n\}$  is Fejér monotone with respect to a subset  $C$  of  $M$  if  $d(x_{n+1}, x) \leq d(x_n, x)$  for all  $x \in C$ .

For a bounded sequence  $\{x_n\}$  in a metric space  $M$ , set

$$r(x, \{x_n\}) = \limsup_{n \rightarrow \infty} d(x, x_n)$$

for all  $x \in M$ .

The asymptotic radius of  $\{x_n\}$  with respect to  $C \subseteq M$  is defined as

$$r(\{x_n\}) = \inf_{x \in C} r(x, \{x_n\}).$$

A point  $y \in C$  is called the asymptotic centre of  $\{x_n\}$  with respect to  $C \subseteq M$  if

$$r(y, \{x_n\}) \leq r(x, \{x_n\}) \text{ for all } x \in C.$$

The set of all asymptotic centres of  $\{x_n\}$  is denoted by  $A(\{x_n\})$ .

A sequence  $\{x_n\}$  in  $M$ , is  $\Delta$ -convergent to  $x \in M$  ( $\Delta - \lim_n x_n = x$ ) if  $x$  is the unique asymptotic centre of  $\{u_n\}$  for every subsequence  $\{u_n\}$  of  $\{x_n\}$ . It has been observed that  $\Delta$ -convergence in metric spaces constitutes an analogue of weak convergence in Hilbert spaces and both coincide in Hilbert spaces.

Let  $(M, d)$  be a metric space. Suppose that there exists a family  $F$  of metric segments such that any two points  $x, y$  in  $M$  are endpoints of a unique metric segment  $[x, y] \in F$  ( $[x, y]$  is an isometric image of the real line interval  $[0, d(x, y)]$ ). We denote by  $z$  the unique point  $\alpha x \oplus (1 - \alpha)y$  of  $[x, y]$  which satisfies

$$d(x, z) = (1 - \alpha)d(x, y) \text{ and } d(z, y) = \alpha d(x, y) \text{ for } \alpha \in I = [0, 1].$$

Such metric spaces are usually called convex metric spaces [18]. A convex metric space  $M$  is hyperbolic if

$$d(\alpha x \oplus (1 - \alpha)y, \alpha z \oplus (1 - \alpha)w) \leq \alpha d(x, z) + (1 - \alpha) d(y, w) \quad (1.3)$$

for all  $x, y, z, w \in M$  and  $\alpha \in I$ .

For  $z = w$ , the hyperbolic inequality reduces to convex structure of Takahashi [23]

$$d(\alpha x \oplus (1 - \alpha)y, z) \leq \alpha d(x, z) + (1 - \alpha)d(y, z).$$

A nonempty subset  $C$  of a hyperbolic space  $M$  is convex if  $\alpha x \oplus (1 - \alpha)y \in C$  for all  $x, y \in C$  and  $\alpha \in I$ . A few examples of nonlinear hyperbolic spaces are Hadamard manifolds [4], the Hilbert open unit ball equipped with the hyperbolic metric [8] and the  $CAT(0)$  spaces [14,15] while normed spaces and their subsets are linear hyperbolic spaces. Throughout this paper, we denote  $\frac{1}{2}x \oplus \frac{1}{2}y$  by  $\frac{x \oplus y}{2}$ .

A hyperbolic space  $M$  is uniformly convex if

$$\delta(r, \varepsilon) = \inf \left\{ 1 - \frac{1}{r} d\left(a, \frac{x \oplus y}{2}\right) : d(a, x) \leq r, d(a, y) \leq r, d(x, y) \geq r\varepsilon \right\} > 0,$$

for any  $a \in M, r > 0$  and  $\varepsilon > 0$ .

Xu [24], extensively used the concept of  $p$ -uniform convexity; its nonlinear version in hyperbolic spaces for  $p = 2$  has been introduced by Khamsi and Khan [13] as under:

For a fixed  $a \in M, r > 0, \varepsilon > 0$ , define

$$\psi(r, \varepsilon) = \inf \left\{ \frac{1}{2}d(a, x)^2 + \frac{1}{2}d(a, y)^2 - d\left(a, \frac{x \oplus y}{2}\right)^2 \right\}$$

where the infimum is taken over all  $x, y \in M$  such that  $d(a, x) \leq r, d(a, y) \leq r$  and  $d(x, y) \geq r\varepsilon$ .

We say that  $M$  is 2-uniformly convex if

$$c_M = \inf \left\{ \frac{\psi(r, \varepsilon)}{r^2 \varepsilon^2} : r > 0, \varepsilon > 0 \right\} > 0.$$

It has been shown in [13] that any  $CAT(0)$  space is 2-uniformly convex hyperbolic space with  $c_M = \frac{1}{4}$ .

From now onwards we assume that  $M$  is a uniformly convex hyperbolic space with the property that for every  $s \geq 0, \varepsilon > 0$ , there exists  $\eta(s, \varepsilon) > 0$  depending on  $s$  and  $\varepsilon$  such that  $\delta(r, \varepsilon) > \eta(s, \varepsilon) > 0$  for any  $r > s$ .

Using the concept of metric segment  $[x, y]$ , we translate (1.2) for nonexpansive mappings in a hyperbolic space as follows:

$$\begin{aligned} x_0 &= x \in C, \\ x_{n+1} &= \alpha_n T\left(\frac{x_n \oplus x_{n+1}}{2}\right) \oplus (1 - \alpha_n)x_n, \end{aligned} \quad (1.4)$$

where  $\{\alpha_n\}$  is the sequence in  $(0, 1)$  satisfying (C1):  $\liminf_{n \rightarrow \infty} \alpha_n > 0$  and (C2):  $\alpha_{n+1}^2 \leq \lambda \alpha_n^2$  for some  $\lambda > 0$ .

The following known results are needed in the sequel.

**Lemma 1.1** ([3]). *Let  $C$  be a nonempty closed subset of a complete metric space  $(M, d)$  and  $\{x_n\}$  be a Fejér monotone with respect to  $C$ . Then  $\{x_n\}$  strongly converges to  $x \in C$  if and only if  $\lim_{n \rightarrow \infty} d(x_n, C) = 0$ .*

**Lemma 1.2** ([6]). *Let  $C$  be a nonempty closed and convex subset of a complete uniformly convex hyperbolic space  $M$ . Then every bounded sequence  $\{y_n\}$  in  $M$  has a unique asymptotic centre with respect to  $C$  that lies in  $C$ .*

**Lemma 1.3** ([10]). *Suppose that  $M$  is a 2-uniformly convex hyperbolic space. Then for any  $\theta \in (0, 1)$ , we have that*

$$d(u, \theta x \oplus (1 - \theta)y)^2 \leq \theta d(u, x)^2 + (1 - \theta)d(u, y)^2 - 4c_M \min\{\theta^2, (1 - \theta)^2\}d(x, y)^2,$$

for all  $u, x, y \in M$  and  $c_M$  is the number as given above.

Our purpose in this paper is to approximate fixed point of nonexpansive mappings using iterative method (1.4) in a 2-uniformly convex hyperbolic spaces. This work provides a unified approach to convergence results in Hilbert spaces, uniformly convex Banach spaces and CAT(0) spaces.

## 2. Convergence in 2-uniformly convex hyperbolic spaces

**Lemma 2.1.** *Let  $C$  be a nonempty convex subset of a complete hyperbolic space  $M$  and  $T : C \rightarrow C$  a nonexpansive mapping. Then the sequence  $\{x_n\}$  in (1.4) is well defined.*

**Proof.** Define  $S : C \rightarrow C$  by

$$Sx = \alpha_0 T\left(\frac{x_0 \oplus x}{2}\right) \oplus (1 - \alpha_0)x_0.$$

With the help of (1.3), we have

$$\begin{aligned} d(Sx, Sy) &= d\left(\alpha_0 T\left(\frac{x_0 \oplus x}{2}\right) \oplus (1 - \alpha_0)x_0, \alpha_0 T\left(\frac{x_0 \oplus y}{2}\right) \oplus (1 - \alpha_0)x_0\right) \\ &\leq \alpha_0 d\left(T\left(\frac{x_0 \oplus x}{2}\right), T\left(\frac{x_0 \oplus y}{2}\right)\right) \\ &\leq \alpha_0 d\left(\frac{x_0 \oplus x}{2}, \frac{x_0 \oplus y}{2}\right) \\ &\leq \frac{\alpha_0}{2} d(x, y). \end{aligned}$$

This gives that  $S$  is a contraction with contraction constant  $\frac{\alpha_0}{2} \in (0, 1)$ . Therefore by Banach contraction principle, there is a unique element  $x_1 \in C$  such that  $x_1 = Sx_1 = \alpha_0 T\left(\frac{x_0 \oplus x_1}{2}\right) \oplus (1 - \alpha_0)x_0$ . Hence  $x_1$  is achieved. Similarly, we can find  $x_2$  and so on. So in general,

$$x_{n+1} = \alpha_n T\left(\frac{x_n \oplus x_{n+1}}{2}\right) \oplus (1 - \alpha_n)x_n. \quad \square$$

**Lemma 2.2.** *Let  $C$  be a nonempty convex subset of a complete 2-uniformly convex hyperbolic space  $M$  and  $T : C \rightarrow C$  a nonexpansive mapping such that  $F(T) \neq \emptyset$ . Then for the sequence  $\{x_n\}$  in (1.4), we have the following: (i)  $\lim_{n \rightarrow \infty} d(x_n, p)$  exists for all  $p \in F(T)$*

- (ii)  $\sum_{n=1}^{\infty} \alpha_n d(x_n, x_{n+1}) < \infty$
- (iii)  $\sum_{n=1}^{\infty} \alpha_n^2 (1 - \alpha_n)^2 d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2 < \infty$ .

**Proof.** Let  $p \in F(T)$ . Applying Lemma 1.3 to (1.4), we have that

$$\begin{aligned}
 d(x_{n+1}, p)^2 &= d\left(\alpha_n T\left(\frac{x_n \oplus x_{n+1}}{2}\right) \oplus (1 - \alpha_n)x_n, p\right)^2 \\
 &\leq \alpha_n d\left(T\left(\frac{x_n \oplus x_{n+1}}{2}\right), p\right)^2 + (1 - \alpha_n)d(x_n, p)^2 \\
 &\quad - 4c_M \min\{\alpha_n^2, (1 - \alpha_n)^2\} d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2 \\
 &\leq \alpha_n d\left(\frac{x_n \oplus x_{n+1}}{2}, p\right)^2 + (1 - \alpha_n)d(x_n, p)^2 \\
 &\quad - 4c_M \alpha_n^2 (1 - \alpha_n)^2 d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2 \\
 &\leq \alpha_n d\left(\frac{1}{2}d(x_n, p)^2 + \frac{1}{2}d(x_{n+1}, p)^2 - \frac{C_M}{4}d(x_n, x_{n+1})^2\right) \\
 &\quad + (1 - \alpha_n)d(x_n, p)^2 - 4c_M \alpha_n^2 (1 - \alpha_n)^2 d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2.
 \end{aligned}$$

That is,

$$\begin{aligned}
 \left(1 - \frac{\alpha_n}{2}\right)d(x_{n+1}, p) &\leq \left(1 - \frac{\alpha_n}{2}\right)d(x_n, p) - \frac{\alpha_n C_M}{4}d(x_n, x_{n+1})^2 \\
 &\quad - 4c_M \alpha_n^2 (1 - \alpha_n)^2 d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2
 \end{aligned}$$

which further implies that

$$\begin{aligned}
 d(x_{n+1}, p) &\leq d(x_n, p) - \frac{\alpha_n C_M}{2(2 - \alpha_n)}d(x_n, x_{n+1}). \\
 &\quad - \frac{8c_M \alpha_n^2 (1 - \alpha_n)^2}{2(2 - \alpha_n)}d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2.
 \end{aligned}$$

The above inequality provides the following three inequalities:

$$d(x_{n+1}, p) \leq d(x_n, p), \tag{2.1}$$

$$\frac{\alpha_n C_M}{2(2 - \alpha_n)}d(x_n, x_{n+1}) \leq d(x_n, p) - d(x_{n+1}, p) \tag{2.2}$$

and

$$\frac{8c_M \alpha_n^2 (1 - \alpha_n)^2}{2(2 - \alpha_n)}d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2 \leq d(x_n, p) - d(x_{n+1}, p). \tag{2.3}$$

From (2.1), it follows that  $\lim_{n \rightarrow \infty} d(x_n, p)$  exists, that is, (i) holds.

Since  $\alpha_n \in (0, 1)$ , therefore  $\alpha_n \leq \frac{\alpha_n}{2(2 - \alpha_n)}$ . Hence (2.2) becomes

$$\alpha_n d(x_n, x_{n+1}) \leq \frac{1}{C_M} [d(x_n, p) - d(x_{n+1}, p)]. \tag{2.4}$$

Let  $m \geq 1$  be any positive integer. Then from (2.4), we have that

$$\sum_{n=1}^m \alpha_n d(x_n, x_{n+1}) \leq \frac{1}{C_M} [d(x_1, p) - d(x_{m+1}, p)] \leq \frac{d(x_1, p)}{C_M}.$$

Let  $m \rightarrow \infty$ . Then

$$\sum_{n=1}^{\infty} \alpha_n d(x_n, x_{n+1}) \leq \frac{d(x_1, p)}{C_M} < \infty.$$

That is,

$$\sum_{n=1}^{\infty} \alpha_n d(x_n, x_{n+1}) < \infty,$$

proving (ii). Similarly, from (2.3), we have

$$\sum_{n=1}^{\infty} \alpha_n^2 (1 - \alpha_n)^2 d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2 < \infty. \quad \square$$

**Lemma 2.3.** *Let  $C$  be a nonempty convex subset of a complete 2-uniformly convex hyperbolic space  $M$  and  $T : C \rightarrow C$  a nonexpansive mapping such that  $F(T) \neq \emptyset$ . Then for the sequence  $\{x_n\}$  in (1.4), we have that  $\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0$ .*

**Proof.** Consider

$$\begin{aligned} d(x_{n+1}, x_{n+2}) &= d\left(\alpha_{n+1} T\left(\frac{x_{n+1} \oplus x_{n+2}}{2}\right) \oplus (1 - \alpha_{n+1})x_{n+1}, x_{n+1}\right) \\ &\leq \alpha_{n+1} d\left(x_{n+1}, T\left(\frac{x_{n+1} \oplus x_{n+2}}{2}\right)\right) \\ &\leq \alpha_{n+1} d\left(x_{n+1}, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) \\ &\quad + \alpha_{n+1} d\left(T\left(\frac{x_n \oplus x_{n+1}}{2}\right), T\left(\frac{x_{n+1} \oplus x_{n+2}}{2}\right)\right) \\ &\leq \alpha_{n+1} d\left(x_{n+1}, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) \\ &\quad + \alpha_{n+1} d\left(\frac{x_n \oplus x_{n+1}}{2}, \frac{x_{n+1} \oplus x_{n+2}}{2}\right) \\ &\leq \alpha_{n+1} (1 - \alpha_n) d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) \\ &\quad + \alpha_{n+1} d\left(\frac{x_n \oplus x_{n+1}}{2}, \frac{x_{n+1} \oplus x_{n+2}}{2}\right) \\ &\leq \alpha_{n+1} (1 - \alpha_n) d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) \\ &\quad + \frac{\alpha_{n+1}}{2} (d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2})). \end{aligned}$$

Therefore

$$\left(1 - \frac{\alpha_{n+1}}{2}\right) d(x_{n+2}, x_{n+1}) \leq \alpha_{n+1} (1 - \alpha_n) d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) + \frac{\alpha_{n+1}}{2} d(x_n, x_{n+1})$$

which further implies that

$$\begin{aligned} d(x_{n+1}, x_{n+2}) &\leq \frac{2\alpha_{n+1}(1-\alpha_n)}{2-\alpha_{n+1}} d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) \\ &\quad + \frac{\alpha_{n+1}}{2-\alpha_{n+1}} d(x_n, x_{n+1}) \\ &\leq 2\alpha_{n+1}(1-\alpha_n) d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) \\ &\quad + \alpha_{n+1} d(x_n, x_{n+1}). \end{aligned}$$

For some  $A > 0, B > 0$  and using the assumption  $\alpha_{n+1}^2 \leq \lambda\alpha_n^2$ , we further derive that

$$\begin{aligned} d(x_{n+1}, x_{n+2})^2 &\leq 4A\alpha_{n+1}^2(1-\alpha_n)^2 d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2 \\ &\quad + B\alpha_{n+1}^2 d(x_n, x_{n+1})^2 \\ &\leq 4A\lambda\alpha_n^2(1-\alpha_n)^2 d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2 \\ &\quad + B\alpha_{n+1}^2 d(x_n, x_{n+1})^2 \\ &\leq 4A\lambda\alpha_n^2(1-\alpha_n)^2 d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2 \\ &\quad + B\lambda\alpha_n^2 d(x_n, x_{n+1})^2 \\ &\leq 4A\lambda\alpha_n^2(1-\alpha_n)^2 d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right)^2 \\ &\quad + B\lambda\alpha_n d(x_n, x_{n+1})^2. \end{aligned}$$

Hence by Lemma 2.2(ii)–(iii), we have that

$$\sum_{n=1}^{\infty} d(x_{n+1}, x_{n+2})^2 < \infty.$$

This in turn implies that

$$\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0. \quad \square \tag{2.5}$$

**Lemma 2.4.** *Let  $C$  be a nonempty closed and convex subset of a complete 2-uniformly convex hyperbolic space  $M$  and  $T : C \rightarrow C$  a nonexpansive mapping such that  $F(T) \neq \emptyset$ . Then for the sequence  $\{x_n\}$  in (1.4), we have that  $\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0$ .*

**Proof.** The condition  $\liminf_{n \rightarrow \infty} \alpha_n > 0$  implies that  $0 < \frac{1}{\alpha_n} \leq \frac{1}{\alpha}$  for sufficiently large  $n$ . The inequality

$$\begin{aligned} d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) &\leq d(x_n, x_{n+1}) + d\left(x_{n+1}, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) \\ &\leq d(x_n, x_{n+1}) + (1-\alpha_n) d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) \end{aligned}$$

implies that

$$d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) \leq \frac{1}{\alpha_n} d(x_n, x_{n+1}) \leq \frac{1}{\alpha} d(x_n, x_{n+1}).$$

By taking  $\limsup_{n \rightarrow \infty}$  on both sides in the above inequality and then appealing to Lemma 2.3, we get that

$$\lim_{n \rightarrow \infty} d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) = 0. \tag{2.6}$$

Finally, the inequality

$$\begin{aligned} d(x_n, T x_n) &\leq d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) + d\left(T\left(\frac{x_n \oplus x_{n+1}}{2}\right), T x_n\right) \\ &\leq d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) + d\left(\frac{x_n \oplus x_{n+1}}{2}, x_n\right) \\ &\leq d\left(x_n, T\left(\frac{x_n \oplus x_{n+1}}{2}\right)\right) + \frac{1}{2} d(x_{n+1}, x_n) \end{aligned}$$

together with (2.5) and (2.6) provides that

$$\lim_{n \rightarrow \infty} d(x_n, T x_n) = 0. \quad \square \tag{2.7}$$

The following concept is needed to establish strong convergence of (1.4).

Let  $f$  be a nondecreasing function on  $[0, \infty)$  with  $f(0) = 0$  and  $f(t) > 0$  for all  $t \in (0, \infty)$ . Then the mapping  $T : C \rightarrow C$  with  $F(T) \neq \phi$ , satisfies condition (A) [21] if

$$d(x, T x) \geq f(d(x, F(T))) \text{ for } x \in C,$$

where  $d(x, F(T)) = \inf\{d(x, y) : y \in F(T)\}$ .

Using condition(A) and Lemma 2.4, we obtain the following strong convergence result.

**Theorem 2.5.** *Let  $C$  be a nonempty closed and convex subset of a complete 2-uniformly convex hyperbolic space  $M$  and  $T : C \rightarrow C$  a nonexpansive mapping such that  $F(T) \neq \phi$ . If the mapping  $T : C \rightarrow C$  satisfies condition(A), then the sequence  $\{x_n\}$  in (1.4), strongly converges to a fixed point of  $T$ .*

**Proof.** By Lemma 2.4,  $\lim_{n \rightarrow \infty} d(x_n, T x_n) = 0$ . Now condition(A) implies that  $\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0$ . Finally, by Lemma 1.1,  $\{x_n\}$  strongly converges to a fixed point of  $T$ .  $\square$

Here are our other strong convergence results.

**Theorem 2.6.** *Let  $C$  be a nonempty closed and convex subset of a complete 2-uniformly convex hyperbolic space  $M$  and  $T : C \rightarrow C$  a nonexpansive mapping such that  $F(T) \neq \phi$ . If  $T$  is semi-compact, then the sequence  $\{x_n\}$  in (1.4) strongly converges to a fixed point of  $T$ .*

**Proof.** By Lemma 2.4, we have that  $\lim_{n \rightarrow \infty} d(x_n, T x_n) = 0$ . Since  $\lim_{n \rightarrow \infty} d(x_n, p)$  exists for each  $p \in F(T)$ ,  $\{x_n\}$  is bounded. As  $\lim_{n \rightarrow \infty} d(x_n, T x_n) = 0$  and  $T$  is semi-compact, so there is a subsequence  $\{x_{n_i}\}$  of  $\{x_n\}$  such that  $x_{n_i} \rightarrow q \in C$  and hence  $T x_{n_i} \rightarrow T q$ . Therefore,  $\lim_{i \rightarrow \infty} d(x_{n_i}, T x_{n_i}) = 0$  implies that  $d(T q, q) = 0$ . That is,  $q \in F(T)$ . Since  $\lim_{n \rightarrow \infty} d(x_n, p)$  exists and  $x_{n_i} \rightarrow q, x_n \rightarrow q$ .  $\square$

**Theorem 2.7.** *Let  $C$  be a nonempty closed and convex subset of a complete 2-uniformly convex hyperbolic space  $M$  and  $T : C \rightarrow C$  a nonexpansive mapping such that  $F(T) \neq \phi$ . If  $T$  is completely continuous, then the sequence  $\{x_n\}$  in (1.4), strongly converges to a fixed point of  $T$ .*

**Proof.** Since  $\{x_n\}$  is bounded and  $T$  is completely continuous,  $\{T x_n\}$  has a convergent subsequence say  $\{T x_{n_i}\}$ . Therefore by (2.7),  $\{x_{n_i}\}$  converges. Let  $\lim_{i \rightarrow \infty} x_{n_i} = v$ . By continuity

of  $T$  and (2.7), we have that  $Tv = v$ . By Lemma 2.2,  $\lim_{n \rightarrow \infty} d(x_n, v)$  exists and so  $\{x_n\}$  strongly converges to  $v$ .  $\square$

We now present our  $\Delta$ -convergence result.

**Theorem 2.8.** *Let  $C$  be a nonempty closed and convex subset of a complete 2-uniformly convex hyperbolic space  $M$  and  $T : C \rightarrow C$  a nonexpansive mapping such that  $F(T) \neq \emptyset$ . Then the sequence  $\{x_n\}$  in (1.4),  $\Delta$ -converges to a fixed point of  $T$ .*

**Proof.** It follows from Lemma 2.1 that  $\{x_n\}$  is bounded in  $C$ . By Lemma 1.2,  $\{x_n\}$  has a unique asymptotic centre, that is,  $A_C(\{x_n\}) = \{y\}$ . Let  $\{w_n\}$  be any subsequence of  $\{x_n\}$  such that  $A_C(\{w_n\}) = \{w\}$ . We claim that  $w \in F(T)$ . By Lemma 2.4, we have that

$$\lim_{n \rightarrow \infty} d(w_n, Tw_n) = 0.$$

The nonexpansive mapping  $T$  satisfies the following inequality:

$$d(w_n, Tw) \leq d(w_n, Tw_n) + d(w_n, w)$$

which further implies that

$$\limsup_{n \rightarrow \infty} d(w_n, Tw) \leq \limsup_{n \rightarrow \infty} d(w_n, Tw_n) + \limsup_{n \rightarrow \infty} d(w_n, w) = \limsup_{n \rightarrow \infty} d(w_n, w).$$

By the uniqueness of asymptotic centre, we have  $Tw = w$ . Therefore  $F(T) \neq \emptyset$ . If  $y \neq w$ , then by the uniqueness of asymptotic centre and the fact that  $\lim_{n \rightarrow \infty} d(x_n, x)$  exists for each  $x \in F(T)$ , we have that

$$\begin{aligned} \limsup_{n \rightarrow \infty} d(w_n, w) &< \limsup_{n \rightarrow \infty} d(w_n, y) \\ &\leq \limsup_{n \rightarrow \infty} d(x_n, y) \\ &< \limsup_{n \rightarrow \infty} d(x_n, w) \\ &= \limsup_{n \rightarrow \infty} d(w_n, w). \end{aligned}$$

This is a contradiction and therefore  $y = w$ . This proves that  $\{x_n\}$ ,  $\Delta$ -converges to  $x \in F(T)$ .  $\square$

**Remark 2.9.** (1) All the results of this paper instantly hold in Hilbert spaces, uniformly convex Banach spaces satisfying Opial property and CAT(0) spaces; (2) The results of Alghamdi et al. [1] are corollaries of our corresponding results; (3) The interested reader is referred to [11] for another notion of  $p$ -uniformly convex metric spaces; (4) The two control conditions: (C1) and (C2) in our algorithm (1.4) are satisfied by the sequence  $\alpha_n = 1 - \frac{1}{n+1}$ .

### 3. Application

We know that  $L^2[0, 1]$  is a Hilbert space and hence it is a 2-uniformly convex hyperbolic space. Suppose that  $h: [0, 1] \rightarrow [0, 1]$  and  $F: [0, 1] \times [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  are continuous functions and  $F$  satisfies the Lipschitz continuity condition, i.e.,

$$|F(t, \lambda, x) - F(t, s, y)| \leq |x - y| \text{ for } t, s \in [0, 1] \text{ and } x, y \in \mathbb{R}.$$

Consider a Fredholm integral equation of the form

$$x(t) = h(t) + \int_0^1 F(t, s, x(s)) ds \quad \text{for } t \in [0, 1]. \tag{3.1}$$

It has been shown in [19] that the solution of Eq. (3.1) exists in  $L^2[0, 1]$ . To find an approximate solution of this equation, we define  $S: L^2[0, 1] \rightarrow L^2[0, 1]$  by

$$Sx(t) = h(t) + \int_0^1 F(t, s, x(s)) ds \quad \text{for } t \in [0, 1].$$

For  $x, y \in L^2 [0, 1]$ , we calculate

$$\begin{aligned} \|Sx - Sy\|^2 &= \int_0^1 |Sx(t) - Sy(t)|^2 dt \\ &= \int_0^1 \left| \int_0^1 (F(t, s, x(s)) - F(t, s, y(s))) ds \right|^2 dt \\ &\leq \int_0^1 \left| \int_0^1 |x(s) - y(s)| ds \right|^2 dt \\ &\leq \int_0^1 |x(s) - y(s)|^2 ds = \|x - y\|^2. \end{aligned}$$

So  $S$  is nonexpansive. For any function  $x_0 \in L^2[0, 1]$ , we define a sequence of functions  $\{x_n\}$  in  $L^2[0, 1]$  by

$$x_{n+1} = \alpha_n S\left(\frac{x_n + x_{n+1}}{2}\right) + (1 - \alpha_n)x_n$$

where  $\alpha_n \in (0, 1)$  such that  $\liminf_{n \rightarrow \infty} \alpha_n > 0$  and  $\alpha_{n+1}^2 \leq \lambda \alpha_n^2$  for some  $\lambda > 0$ . Now by [Theorem 2.8](#),  $\{x_n\}$  weakly converges to the fixed point of  $S$  which is a solution of Eq. (3.1).

## References

- [1] M.A. Alghamdi, M.A. Alghamdi, N. Shahzad, H.K. Xu, The implicit midpoint rule for nonexpansive mappings, *Fixed Point Theory Appl.* 2014 (2014) 96.
- [2] G. Bader, P. Deuffhard, A semi-implicit mid-point rule for stiff systems of ordinary differential equations, *Numer. Math.* 41 (1983) 373–398.
- [3] H.H. Bauschke, P.L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, Springer-Verlag, New York, 2011.
- [4] H. Busemann, Spaces with non-positive curvature, *Acta. Math.* 80 (1948) 259–310.
- [5] P. Deuffhard, Recent progress in extrapolation methods for ordinary differential equations, *SIAM Rev.* 27 (4) (1985) 505–535.
- [6] H. Fukhar-ud din, Existence and approximation of fixed points in convex metric spaces, *Carpathian J. Math.* 30 (2014) 175–185.
- [7] H. Fukhar-ud din, A.R. Khan, A. Kalsoom, M.A.A. Khan, One-step implicit algorithm for two finite families of nonexpansive maps in hyperbolic spaces, *J. Adv. Math. Stud.* 6 (2013) 73–81.
- [8] K. Goebel, S. Reich, *Uniform convexity, and hyperbolic geometry nonexpansive mappings*, Series of Monographs and Textbooks in Pure and Applied Mathematics, Vol. 83, Dekker, New York, 1984.
- [9] B. Halpern, Fixed points of nonexpanding maps, *Bull. Amer. Math. Soc.* 73 (1967) 591–597.
- [10] B.A. Ibn Dehaish, M.A. Khamsi, A.R. Khan, Mann iteration process for asymptotic pointwise nonexpansive mappings in metric spaces, *J. Math. Anal. Appl.* 397 (2013) 861–868.
- [11] C. Izchukwu, G.C. Ugwunnadi, O.T. Mewomo, A.R. Khan, M. Abbas, Proximal-type algorithms for split minimization problem in P-uniformly convex metric spaces, *Numer. Algorithms* 2018 <http://dx.doi.org/10.1007/s11075-018-0633-9>.
- [12] K.R. Kazmi, S.H. Rizvi, Implicit iterative method for approximating a common solution of split equilibrium problem and fixed point problem for a nonexpansive semigroup, *Arab. J. Math. Sci.* 20 (2014) 57–75.

- [13] M.A. Khamsi, A.R. Khan, Inequalities in metric spaces with applications, *Nonlinear Anal.* 74 (2011) 4036–4045.
- [14] W.A. Kirk, A fixed point theorem in CAT(0)spaces and r-trees, *Fixed Point Theory Appl.* 4 (2004) 309–316.
- [15] L. Leustean, A quadratic rate of asymptotic regularity for CAT(0)-spaces, *J. Math. Anal. Appl.* 325 (2007) 386–399.
- [16] G. López, V. Martín-Márquez, H.K. Xu, Iterative algorithms for the multiple-sets split feasibility problem, in: Y. Censor, M. Jiang, G. Wang (Eds.), *Biomedical Mathematics: Promising Directions in Imaging, Therapy Planning and Inverse Problems*, Medical Physics Publishing, Madison, 2010, pp. 243–279.
- [17] W.R. Mann, Mean value methods in iteration, *Proc. Amer. Math. Soc.* 4 (1953) 506–510.
- [18] K. Menger, *Untersuchungen ü*, *Math. Ann.* 100 (1928) 75–163.
- [19] J.J. Nieto, H.K. Xu, Solvability of nonlinear volterra and fredholm equations in weighted spaces, *Nonlinear Anal.* 24 (1995) 1289–1297.
- [20] C. Schneider, Analysis of the linearly implicit mid-point rule for differential-algebra equations, *Electron. Trans. Numer. Anal.* 1 (1993) 1–10.
- [21] H.F. Senter, W.G. Dotson, Approximatig fixed points of nonexpansive mappings, *Proc. Amer. Math. Soc.* 44 (1974) 375–380.
- [22] S. Somalia, Implicit midpoint rule to the nonlinear degenerate boundary value problems, *Int. J. Comput. Math.* 79 (2002) 327–332.
- [23] W. Takahashi, A convexity in metric spaces and nonexpansive mappings, *Kodai Math. Sem. Rep.* 22 (1970) 142–149.
- [24] H.K. Xu, Inequalities in Banach spaces with applications, *Nonlinear Anal.* 16 (1991) 1127–1138.
- [25] H.K. Xu, R.G. Ori, An implicit iteration process for nonexpansive mappings, *Numer. Funct. Anal. Optim.* 22 (2001) 767–773.

**Corresponding author**

H. Fukhar-ud-din can be contacted at: [hfdin@kfupm.edu.sa](mailto:hfdin@kfupm.edu.sa)

For instructions on how to order reprints of this article, please visit our website:

[www.emeraldgrouppublishing.com/licensing/reprints.htm](http://www.emeraldgrouppublishing.com/licensing/reprints.htm)

Or contact us for further details: [permissions@emeraldinsight.com](mailto:permissions@emeraldinsight.com)