

Fixed point theorems for Geraghty-type mappings applied to solving nonlinear Volterra-Fredholm integral equations in modular G -metric spaces

Godwin Amechi Okeke and Daniel Francis
*Department of Mathematics, School of Physical Sciences,
Federal University of Technology, Owerri, Nigeria*

Abstract

Purpose – The authors prove the existence and uniqueness of fixed point of mappings satisfying Geraghty-type contractions in the setting of preordered modular G -metric spaces. The authors apply the results in solving nonlinear Volterra-Fredholm-type integral equations. The results extend generalize compliment and include several known results as special cases.

Design/methodology/approach – The results of this paper are theoretical and analytical in nature.

Findings – The authors prove the existence and uniqueness of fixed point of mappings satisfying Geraghty-type contractions in the setting of preordered modular G -metric spaces. apply the results in solving nonlinear Volterra-Fredholm-type integral equations. The results extend, generalize, compliment and include several known results as special cases.

Research limitations/implications – The results are theoretical and analytical.

Practical implications – The results were applied to solving nonlinear integral equations.

Social implications – The results has several social applications.

Originality/value – The results of this paper are new.

Keywords Fixed point, Preordered, Modular G -metric spaces, Contractive mapping, Existence and uniqueness, Nonlinear Volterra-Fredholm integral equations

Paper type Research paper

1. Introduction

In 1973, Geraghty [1] introduced an interesting generalization of Banach contraction mapping principle using the concept of class $\mathcal{S}$ of functions, that is $\alpha : \mathbb{R}_+ \rightarrow [0, 1)$ with the condition that $\alpha(t_n) \rightarrow 1 \Rightarrow t_n \rightarrow 0$ where $\mathbb{R}_+$ is the set of all nonnegative real numbers and $t \in \mathbb{R}_+$ for all $n \in \mathbb{N}$. In 2012, Gordji *et al.* [2] proved some fixed point theorems for generalized Geraghty contraction in partially ordered complete metric spaces. Bhaskar and

JEL Classification — 47H09, 47H10, 06A75.

© Godwin Amechi Okeke and Daniel Francis. Published in *Arab Journal of Mathematical Sciences*. Published by Emerald Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licenses/by/4.0/legalcode>

The authors wish to thank the editor and the referees for their comments and suggestions. This paper was completed while the first author was visiting the Abdus Salam School of Mathematical Sciences (ASSMS), Government College University Lahore, Pakistan as a postdoctoral fellow.

Authors Contributions: All authors contributed equally to the writing of this paper.

Conflicts of Interest: The authors declare no conflict of interests.



Lakshmikantham [3] proved a fixed point theorem for a mixed monotone mapping in a metric space endowed with partial order, using a weak contractivity type of assumption. Yolacan [4] established some new fixed point theorems in 0-complete ordered partial metric spaces. He also remarked on coupled generalized Banach contraction mapping. Faraji *et al.* [5] extended some fixed point theorems for Geraghty contractive mappings in b-complete b-metric spaces.

Furthermore, Gupta *et al.* [6], established some fixed point theorems in an ordered complete metric space using distance function. Chaipunya *et al.* [7] proved some fixed point theorems of Geraghty-type contractions concerning the existence and uniqueness of fixed points under the setting of modular metric spaces which also generalized the results in Gordji *et al.* [2] under the influence of a modular metric space.

Geraghty-type contractive mappings in metric spaces was generalized to the concept of preordered G-metric spaces in [8] and the authors in [8] obtained unique fixed point results. Furthermore, other interesting fixed point results in G-metric spaces can be found in [9] and the references therein.

In 2010, an essential study by Chistyakov [10] introduced an aspect of metric called modular metric spaces or parameterized metric space with the time parameter λ (say) and his purpose was to define the notion of a modular on an arbitrary set, develop the theory of metric spaces generated by modulars, called modular metric spaces and, on the basis of it, defined new metric spaces of (multi-valued) functions of bounded generalized variation of a real variable with values in metric semigroups and abstract convex cones.

In the same year, Chistyakov [11], as an application presented an exhausting description of Lipschitz continuous and some other classes of superposition (or Nemytskii) operators, acting in these modular metric spaces. He developed the theory of metric spaces generated by modulars and extended the results given by Nakano [12], Musielak and Orlicz [13], Musielak [14] to modular metric spaces. Modular spaces are extensions of Lebesgue, Riesz and Orlicz spaces of integrable functions.

Modular theories on linear spaces can be found in Nakano [12, 15], where he developed a spectral theory in semi-ordered linear spaces (vector lattices) and established the integral representation for projections acting in this modular space.

Nakano [12] established some modulars on real linear spaces which are convex functionals. Non-convex modulars and the corresponding modular linear spaces were constructed by Musielak and Orlicz [13]. Orlicz spaces and modular linear spaces have already become classical tools in modern nonlinear functional analysis.

Furthermore, the development of theory of metric spaces generated by modulars, called modular metric spaces attracted the attention of several mathematicians (see, e.g. [16–19]).

Okeke *et al.* [20] established some convergence results for three multi-valued ρ -quasi-nonexpansive mappings using a three step iterative scheme. Moreover, these fixed point results are applicable to nonlinear integral and differential equations see [19, 21–26] and the references therein, while [7] deals with application to partial differential equation in modular metric spaces.

In 2013, Azadifar *et al.* [27] introduced the notion of modular G-metric space and proved some fixed point theorems for contractive mappings defined on modular G-metric spaces. Based on definitions given in [27], we intend to extend the fixed point theorems obtained in [7] to preordered modular G-metric spaces in this paper. Furthermore, we prove some fixed point theorems for Geraghty-type contraction mappings in the setting of preordered modular G-metric spaces. We apply our results in proving the existence of a unique solution for a system of nonlinear Volterra-Fredholm integral equations in modular G-metric spaces, $X_{\omega G}$.

2. Preliminaries

We begin this section with the following results and definitions which will be useful in this paper.

Theorem 2.1.

[28] If $\{a_n\}_{n \in \mathbb{N}}$, $\{b_n\}_{n \in \mathbb{N}}$, $\{c_n\}_{n \in \mathbb{N}}$ are three sequences in $\mathbb{R}$ such that

- (1) $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = \ell$,
 - (2) for some positive integer N , $a_n \leq c_n \leq b_n$ for all $n \geq N$.
- Then $\lim_{n \rightarrow \infty} c_n = \ell$.

Definition 2.1. [29] A preorder set X is a relation $\leq$ that is both,

- (1) transitive i.e; $x \leq y$ and $y \leq z$ implies $x \leq z$ and,
- (2) reflexive i.e; $x \leq x$.

A preordered set is a pair $(X, \leq)$ consisting of a set X and a preorder $\leq$ on X .

Remark 2.1. If a preorder $\leq$ is antisymmetric i.e; $x \leq y$ and $y \leq x$ implies $x = y$, then $\leq$ is called a partial order.

Definition 2.2. [1] Let $\mathcal{S}$ be the family of all Geraghty functions, that is functions $\alpha : [0, \infty) \rightarrow [0, 1)$ satisfying the condition $\{\alpha(t_n)\} \rightarrow 1 \Rightarrow \{t_n\} \rightarrow 0$.

For the rest of this paper, we denote the the class of all Geraghty functions by $\mathcal{S}_{Ger}$. Such Geraghty class was discussed in [7].

Definition 2.3. [7] Let $\mathcal{S}$ be the family of all Geraghty functions, that is functions $\beta_i : \mathbb{R}_+ \cup \{\infty\} \rightarrow [0, 1)$ satisfying the condition $\beta_i(t_k) \rightarrow \frac{1}{n} \Rightarrow \{t_k\} \rightarrow 0$ for all i .

Definition 2.4. [7] Let Ψ be the class of functions $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that the following conditions hold;

- (1) ψ is decreasing,
- (2) ψ is continuous,
- (3) $\psi(t) = 0$ if and only if $t = 0$.

Extension of Definition 2.2 above is as follows:

Definition 2.5. [7] Let $\bar{\Psi}$ be the class of functions $\psi : \mathbb{R}_+ \cup \{\infty\} \rightarrow \mathbb{R}_+ \cup \{\infty\}$ such that the following conditions hold;

- (1) ψ is subadditive,
- (2) $\psi(t)$ is finite for $0 < t < \infty$,
- (3) $\psi|_{\mathbb{R}_+} \in \bar{\Psi}$.

Definition 2.6. [27] Let X be a nonempty set, and let $\omega^G : (0, \infty) \times X \times X \times X \rightarrow [0, \infty]$ be a function satisfying;

- (1) $\omega_\lambda^G(x, y, z) = 0$ for all $x, y \in X$ and $\lambda > 0$ if $x = y = z$,
- (2) $\omega_\lambda^G(x, x, y) > 0$ for all $x, y \in X$ and $\lambda > 0$ with $x \neq y$,
- (3) $\omega_\lambda^G(x, x, y) \leq \omega_\lambda^G(x, y, z)$ for all $x, y, z \in X$ and $\lambda > 0$ with $z \neq y$,
- (4) $\omega_\lambda^G(x, y, z) = \omega_\lambda^G(x, z, y) = \omega_\lambda^G(y, z, x) = \dots$ for all $\lambda > 0$ (symmetry in all three variables),

(5) $\omega_{\lambda+\mu}^G(x, y, z) \leq \omega_{\lambda}^G(x, a, a) + \omega_{\mu}^G(a, y, z)$, for all $x, y, z, a \in X$ and $\lambda, \mu > 0$, then the function ω_{λ}^G is called a modular G -metric on X .

Remarks 2.1.

- (1) The pair (X, ω^G) is called a modular G -metric space, and without any confusion we will take X_{ω^G} as a modular G -metric space.
From condition (5), if ω^G is convex, then we have a strong form as,
- (2) $\omega_{\lambda+\mu}^G(x, y, z) \leq \omega_{\frac{\lambda}{\lambda+\mu}}^G(x, a, a) + \omega_{\frac{\mu}{\lambda+\mu}}^G(a, y, z)$,
- (3) If $x = a$, then (5) above becomes $\omega_{\lambda+\mu}^G(a, y, z) \leq \omega_{\mu}^G(a, y, z)$,
- (4) Condition (5) is called rectangle inequality.

Definition 2.7. [27] Let (X, ω^G) be a modular G -metric space. The sequence $\{x_n\}_{n \in \mathbb{N}}$ in X is ω^G -convergent to x , if it converges to x in the topology $\tau(\omega_{\lambda}^G)$.

A function $T : X_{\omega^G} \rightarrow X_{\omega^G}$ at $x \in X_{\omega^G}$ is called ω^G -continuous if $\omega_{\lambda}^G(x_n, x, x) \rightarrow 0$ then $\omega_{\lambda}^G(Tx_n, Tx, Tx) \rightarrow 0$, for all $\lambda > 0$.

Remark 2.2. $\{x_n\}_{n \in \mathbb{N}}$ modular G -converges to x as $n \rightarrow \infty$, if $\lim_{n \rightarrow \infty} \omega_{\lambda}^G(x_n, x_m, x) = 0$. That is for all $\epsilon > 0$ there exists $n_0 \in \mathbb{N}$ such that $\omega_{\lambda}^G(x_n, x_m, x) < \epsilon$ for all $n, m \geq n_0$. Here we say that x is modular G -limit of $\{x_n\}_{n \in \mathbb{N}}$.

Definition 2.8. [27] Let (X, ω^G) be a modular G -metric space, then $\{x_n\}_{n \in \mathbb{N}} \subseteq X_{\omega^G}$ is said to be ω^G -Cauchy if for every $\epsilon > 0$, there exists $n_{\epsilon} \in \mathbb{N}$ such that $\omega_{\lambda}^G(x_n, x_m, x_l) < \epsilon$ for all $n, m, l \geq n_{\epsilon}$ and $\lambda > 0$.

A modular G -metric space X_{ω^G} is said to be ω^G -complete if every ω^G -Cauchy sequence in X_{ω^G} is ω^G -convergent in X_{ω^G} .

Proposition 2.2. [27] Let (X, ω^G) be a modular G -metric space, for any $x, y, x, a \in X$, it follows that:

- (1) If $\omega_{\lambda}^G(x, y, z) = 0$ for all $\lambda > 0$, then $x = y = z$.
- (2) $\omega_{\lambda}^G(x, y, z) \leq \omega_{\lambda/2}^G(x, x, y) + \omega_{\lambda/2}^G(x, x, z)$ for all $\lambda > 0$.
- (3) $\omega_{\lambda}^G(x, y, y) \leq 2\omega_{\lambda/2}^G(x, x, y)$ for all $\lambda > 0$.
- (4) $\omega_{\lambda}^G(x, y, z) \leq \omega_{\lambda/2}^G(x, a, z) + \omega_{\lambda/2}^G(a, y, z)$ for all $\lambda > 0$.
- (5) $\omega_{\lambda}^G(x, y, z) \leq \frac{2}{3}(\omega_{\lambda/2}^G(x, y, a) + \omega_{\lambda/2}^G(x, a, z) + \omega_{\lambda/2}^G(a, y, z))$ for all $\lambda > 0$.
- (6) $\omega_{\lambda}^G(x, y, z) \leq \omega_{\lambda/2}^G(x, a, a) + \omega_{\lambda/2}^G(y, a, a) + \omega_{\lambda/2}^G(z, a, a)$ for all $\lambda > 0$.

Proposition 2.3. [27] Let (X, ω^G) be a modular G -metric space and $\{x_n\}_{n \in \mathbb{N}}$ be a sequence in X_{ω^G} . Then the following are equivalent:

- (1) $\{x_n\}_{n \in \mathbb{N}}$ is ω^G -convergent to x ,
- (2) $\omega_{\lambda}^G(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$, i.e; $\{x_n\}_{n \in \mathbb{N}}$ converges to x relative to modular metric $\omega_{\lambda}^G(\cdot)$,
- (3) $\omega_{\lambda}^G(x_n, x_n, x) \rightarrow 0$ as $n \rightarrow \infty$ for all $\lambda > 0$,
- (4) $\omega_{\lambda}^G(x_n, x, x) \rightarrow 0$ as $n \rightarrow \infty$ for all $\lambda > 0$,
- (5) $\omega_{\lambda}^G(x_m, x_n, x) \rightarrow 0$ as $m, n \rightarrow \infty$ for all $\lambda > 0$.

We give the following definition which will be useful in our results.

Definition 2.9. An ordered modular G -metric space is a triple $(X, \omega^G, \leq)$ where (X, ω) is a modular metric space and $\leq$ is a partial order on X_{ω^G} . If $\leq$ is a preorder on X_{ω^G} , then $(X, \omega^G, \leq)$ is a preordered modular G -metric space.

3. Main results

Theorem 3.1. Let (X, ω^G) be a complete modular G -metric space with a preorder, $\leq$ and a nondecreasing self-mapping $T : X_{\omega^G} \rightarrow X_{\omega^G}$ on X_{ω^G} such that for each $\lambda > 0$, there is $\nu(\lambda) \in [0, \lambda)$ such that the following conditions hold:

(1)

$$\begin{aligned} \psi\left(\omega_\lambda^G(Tx, Ty, Ty)\right) &\leq \alpha\left(\psi\left(\omega_\lambda^G(x, y, y)\right)\right)\psi\left(\omega_{\lambda+\nu(\lambda)}^G(x, y, y)\right) \\ &+ \beta\left(\psi\left(\omega_\lambda^G(x, y, y)\right)\right)\psi\left(\omega_\lambda^G(x, Tx, Tx)\right) + \gamma\left(\psi\left(\omega_\lambda^G(x, y, y)\right)\right)\psi\left(\omega_\lambda^G(y, Ty, Ty)\right), \end{aligned} \quad (3.1)$$

where $\psi \in \bar{\Psi}$ and $\{\alpha, \beta, \gamma\} \in \mathcal{S}_{Ger}$ with $\alpha(t) + 2 \max\{\sup_{t \geq 0} \beta(t), \sup_{t \geq 0} \gamma(t)\} < 1$, and distinct $x, y \in X_{\omega^G}$. Assuming that if a nondecreasing sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to x^* , then $x_n \leq x^*$ for each $n \in \mathbb{N}$,

- (2) if ψ is subadditive and for any $x, y \in X_{\omega^G}$, there exists $z \in X_{\omega^G}$ with $z \leq Tz$ and $\omega_\lambda^G(z, Tz, Tz)$ is finite for all $\lambda > 0$ such that z is comparable to both x and y . Then T has a fixed point $u \in X_{\omega^G}$ and the sequence define by $\{T^n x_0\}_{n \geq 1}$ converges to u . Moreover, the fixed point of T is unique.

Proof. Let $x_0 \in X_{\omega^G}$ be such that $x_0 \leq Tx_0$ and let $x_n = Tx_{n-1} = T^n x_0$ for all $n \in \mathbb{N}$. Regarding that T is nondecreasing mapping, we have that $x_0 \leq Tx_0 = x_1$, which implies that $x_1 = Tx_0 \leq Tx_1 = x_2$. Inductively, we have

$$x_0 \leq x_1 \leq x_2 \leq \dots \leq x_{n-1} \leq x_n \leq x_{n+1} \leq \dots \quad (3.2)$$

Assume that there exists $n_0 \in \mathbb{N}$ such that $x_{n_0} = x_{n_0+1}$. Since $x_{n_0} = x_{n_0+1} = Tx_{n_0}$, then x_{n_0} is the fixed point of T . Now suppose that $x_n \not\leq x_{n+1}$ for all $n \in \mathbb{N}$, thus by inequality (3.2), we have that

$$x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n < x_{n+1} < \dots \quad (3.3)$$

Now for each $\lambda > 0$, and $x_0 < Tx_0$ for all $n \in \mathbb{N}$ implies that $\omega_\lambda^G(x_0, Tx_0, Tx_0) > 0$. Again, let $x_0 \in X_{\omega^G}$ such that $\omega_\lambda^G(x_0, Tx_0, Tx_0) < \infty \forall \lambda > 0$.

First, we show that for all $n \in \mathbb{N}$, the sequence $\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0) = 0$ for all $\lambda > 0$, as $n \rightarrow \infty$. Assume that, for each $n \in \mathbb{N}$, there exists $\lambda_n > 0$ such that $\omega_{\lambda_n}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0) \neq 0$. Otherwise the proof is complete. Suppose not, for each $n \geq 1$, if $0 < \lambda < \lambda_n$, then we have that $\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0) \neq 0$. Since $T^n x_0 \leq T^{n+1} x_0$, from inequality (3.1) we can see that $\psi\left(\omega_{\lambda_n}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)\right) \leq \psi\left(\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)\right) = \psi\left(\omega_\lambda^G(TT^{n-1} x_0, TT^n x_0, TT^n x_0)\right)$. Take $x = T^{n-1} x_0$ and $y = T^n x_0$, then inequality (3.1) becomes;

$$\begin{aligned}
 \psi\left(\omega_{\lambda n}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)\right) &\leq \psi\left(\omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)\right) \\
 &\leq \alpha\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) \times \psi\left(\omega_{\lambda+\nu(\lambda)}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right) \\
 &+ \beta\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) \times \psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right) \\
 &+ \gamma\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) \times \psi\left(\omega_{\lambda}^G(T^n x_0, T^n x_0, T^n x_0)\right) \\
 &= \alpha\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) \times \psi\left(\omega_{\lambda+\nu(\lambda)}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right) \\
 &+ \beta\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) \times \psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right) \\
 &+ \gamma\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) \times \psi\left(\omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)\right) \\
 &\leq \alpha\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) \times \psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right) \\
 &+ \beta\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) \times \psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right) \\
 &+ \gamma\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) \times \psi\left(\omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)\right),
 \end{aligned} \tag{3.4}$$

for which we have that

$$\begin{aligned}
 \psi\left(\omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)\right) &\leq \delta \psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right) \\
 &\leq \psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right) \\
 &\vdots \\
 &\leq \psi\left(\omega_{\lambda}^G(x_0, T x_0, T x_0)\right) \\
 &< \infty,
 \end{aligned} \tag{3.5}$$

where

$$\delta := \frac{\alpha\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) + \beta\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right)}{1 - \gamma\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right)}. \tag{3.6}$$

Therefore, $\{\psi(\omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0))\}_{n \geq 1}$ is nonincreasing and bounded below, so the sequence $\{\psi(\omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0))\}_{n \geq 1}$ converges to some real number $k \geq 0$. Assume $k > 0$, we can see clearly that by using inequality 3, inequality 3 becomes

$$\begin{aligned}
 \psi\left(\omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)\right) &\leq \left(\alpha\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right)\right. \\
 &\quad \left.+ \beta\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) + \gamma\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right)\right)
 \end{aligned} \tag{3.7}$$

as $n \rightarrow \infty$, we get

$$\begin{aligned}
 1 &\leq \liminf_{n \rightarrow \infty} \left(\alpha\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right) + \beta\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right)\right. \\
 &\quad \left.+ \gamma\left(\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right)\right)\right)
 \end{aligned} \tag{3.8}$$

So, we have that

$$\psi\left(\omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0)\right) = 0, \tag{3.9}$$

hence

$$\lim_{n \rightarrow \infty} \omega_{\lambda}^G(T^{n-1} x_0, T^n x_0, T^n x_0) = 0 \tag{3.10}$$

for all $\lambda > 0$, which is a contradiction to our assumption. Therefore,

$$\lim_{n \rightarrow \infty} \psi(\omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)) = 0, \quad (3.11)$$

so,

$$\lim_{n \rightarrow \infty} \omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0) = 0 \quad (3.12)$$

220

for all $\lambda > 0$. This shows that $\omega_{\lambda}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0) = 0$ for all $\lambda > 0, n \geq 1$.

Next, we show that $\{T^n x_0\}_{n \geq 1}$ is a modular G -Cauchy sequence. Suppose, if possible that $\{T^n x_0\}_{n \geq 1}$ not a modular G -Cauchy sequence, then there exists real numbers, $\lambda_0 > 0, \epsilon > 0$ and also there exists two subsequences $\{T^{n_k} x_0\}_{k \geq 1}$ and $\{T^{m_k} x_0\}_{k \geq 1}$ of the sequence $\{T^n x_0\}_{n \geq 1}$ such that, for $n_k > m_k > k$, we have that $\omega_{\lambda_0}^G(T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0) \geq \epsilon$, but $\omega_{\lambda_0}^G(T^{m_k} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) < \epsilon$. Now, since $T^{m_k} x_0 \leq T^{n_k} x_0$, we have that $\epsilon \leq \omega_{\lambda_0}^G(T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0)$ which implies that $\psi(\epsilon) \leq \psi(\omega_{\lambda_0}^G(T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0)) = \psi(\omega_{\lambda_0}^G(TT^{m_k-1} x_0, TT^{n_k-1} x_0, TT^{n_k-1} x_0))$. Set $x = T^{m_k-1} x_0$ and $y = T^{n_k-1} x_0$ into inequality (3.1), then we have

$$\begin{aligned} \psi(\epsilon) &\leq \psi\left(\omega_{\lambda_0}^G(T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0)\right) \leq \alpha\left(\psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right)\right) \\ &\times \psi\left(\omega_{\lambda_0+\nu(\lambda_0)}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right) + \beta\left(\psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right)\right) \\ &\times \psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, TT^{m_k-1} x_0, TT^{m_k-1} x_0)\right) + \gamma\left(\psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right)\right) \\ &\times \psi\left(\omega_{\lambda_0}^G(T^{n_k-1} x_0, TT^{n_k-1} x_0, TT^{n_k-1} x_0)\right) = \alpha\left(\psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right)\right) \\ &\times \psi\left(\omega_{\lambda_0+\nu(\lambda_0)}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right) + \beta\left(\psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right)\right) \\ &\times \psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0)\right) + \gamma\left(\psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right)\right) \\ &\times \psi\left(\omega_{\lambda_0}^G(T^{n_k-1} x_0, T^{n_k} x_0, T^{n_k} x_0)\right) \leq \alpha\left(\psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right)\right) \\ &+ \psi\left(\omega_{\lambda_0}^G(T^{m_k} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right) + \beta\left(\psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right)\right) \\ &\times \psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0)\right) + \gamma\left(\psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right)\right) \\ &\times \psi\left(\omega_{\lambda_0}^G(T^{n_k-1} x_0, T^{n_k} x_0, T^{n_k} x_0)\right) \leq \psi\left(\omega_{\nu(\lambda_0)}^G(T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0)\right) \\ &+ \psi\left(\omega_{\lambda_0}^G(T^{m_k} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0)\right) + \psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0)\right) \\ &+ \psi\left(\omega_{\lambda_0}^G(T^{n_k-1} x_0, T^{n_k} x_0, T^{n_k} x_0)\right) \leq \psi\left(\omega_{\nu(\lambda_0)}^G(T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0)\right) \\ &+ \psi(\epsilon) + \psi\left(\omega_{\lambda_0}^G(T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0)\right) + \psi\left(\omega_{\lambda_0}^G(T^{n_k-1} x_0, T^{n_k} x_0, T^{n_k} x_0)\right), \end{aligned} \quad (3.13)$$

as $k \rightarrow \infty$, we obtain

$$\psi(\epsilon) \leq \lim_{k \rightarrow \infty} \psi\left(\omega_{\lambda_0}^G(T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0)\right) \leq \psi(\epsilon), \quad (3.14)$$

so that

$$\lim_{k \rightarrow \infty} \psi \left(\omega_{\lambda_0}^G (T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0) \right) = \psi(\epsilon). \quad (3.15)$$

Hence

$$\lim_{k \rightarrow \infty} \omega_{\lambda_0}^G (T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0) = \epsilon. \quad (3.16)$$

Again, using condition 5 of [Definition 2.6](#), we get

$$\begin{aligned} \psi \left(\omega_{\lambda_0}^G (T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0) \right) &\leq \alpha \left(\psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) \right) \right) \\ &\times \psi \left(\omega_{\lambda_0 + \nu(\lambda_0)}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) \right) + \beta \left(\psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{m_k-1} x_0) \right) \right) \\ &\times \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0) \right) + \gamma \left(\psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{m_k-1} x_0) \right) \right) \\ &\times \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k} x_0, T^{n_k} x_0) \right) \leq \psi \left(\omega_{\lambda_0 + \nu(\lambda_0)}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) \right) \\ &+ \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0) \right) + \psi \left(\omega_{\lambda_0}^G (T^{n_k-1} x_0, T^{n_k} x_0, T^{n_k} x_0) \right) \\ &\leq \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) \right) + \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0) \right) \\ &+ \psi \left(\omega_{\lambda_0}^G (T^{n_k-1} x_0, T^{n_k} x_0, T^{n_k} x_0) \right) \leq \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) \right) \\ &+ \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0) \right) + \psi \left(\omega_{\frac{\lambda_0}{2}}^G (T^{n_k-1} x_0, T^{m_k} x_0, T^{n_k} x_0) \right) \\ &+ \psi \left(\omega_{\frac{\lambda_0}{2}}^G (T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0) \right) \leq \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) \right) \\ &+ \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0) \right) + \psi \left(\omega_{\lambda_0}^G (T^{n_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0) \right) \\ &+ \psi \left(\omega_{\lambda_0}^G (T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0) \right) \leq \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) \right) \\ &+ \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0) \right) + \psi \left(\omega_{\lambda_0}^G (T^{n_k-1} x_0, T^{m_k} x_0, T^{m_k} x_0) \right) \\ &+ \psi \left(\omega_{\lambda_0}^G (T^{m_k} x_0, T^{n_k} x_0, T^{n_k} x_0) \right), \end{aligned} \quad (3.17)$$

as $k \rightarrow \infty$, we have

$$\psi(\epsilon) \leq \lim_{k \rightarrow \infty} \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) \right) \leq \psi(\epsilon), \quad (3.18)$$

so that

$$\lim_{k \rightarrow \infty} \psi \left(\omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) \right) = \psi(\epsilon). \quad (3.19)$$

Hence

$$\lim_{k \rightarrow \infty} \omega_{\lambda_0}^G (T^{m_k-1} x_0, T^{n_k-1} x_0, T^{n_k-1} x_0) = \epsilon. \quad (3.20)$$

Thus, it follows that

$$1 \leq \liminf_{k \rightarrow \infty} \left(\alpha \left(\psi \left(\omega_{\lambda_0}^G (T^{m_k-1}x_0, T^{n_k-1}x_0, T^{n_k-1}x_0) \right) \right) \right). \quad (3.21)$$

Therefore, we conclude that

$$\lim_{k \rightarrow \infty} \omega_{\lambda_0}^G (T^{m_k-1}x_0, T^{n_k-1}x_0, T^{n_k-1}x_0) = 0 \quad \forall \lambda > 0. \quad (3.22)$$

This is a contradiction. Therefore, it follows that $\{T^n x_0\}_{n \geq 1}$ is a modular G -Cauchy sequence in X_{ω^G} . Since X_{ω^G} is complete modular G -metric space, there exists $u \in X_{\omega^G}$ such that $T^n x_0 \rightarrow u \in X_{\omega^G}$. Now we show that u is a fixed point of T for any arbitrary $\lambda > 0$, using condition 5 of Definition 2.6 and inequality (3.1), we have that

$$\begin{aligned} & \psi(\omega_{\lambda}^G(T^n x_0, Tu, Tu)) \leq \psi\left(\omega_{\frac{\lambda}{2}}^G(T^{n+1}x_0, Tu, Tu)\right) \\ & + \psi\left(\omega_{\frac{\lambda}{2}}^G(T^n x_0, T^{n+1}x_0, T^{n+1}x_0)\right) \\ & = \psi\left(\omega_{\lambda/2}^G(T^{n+1}x_0, Tu, Tu)\right) \\ & + \psi\left(\omega_{\lambda/2}^G(T^n x_0, T^{n+1}x_0, T^{n+1}x_0)\right) \\ & \leq \psi(\omega_{\lambda}^G(T^{n+1}x_0, Tu, Tu)) \\ & + \psi(\omega_{\lambda}^G(T^n x_0, T^{n+1}x_0, T^{n+1}x_0)) \\ & \leq \alpha(\psi(\omega_{\lambda}^G(T^n x_0, u, u)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n x_0, u, u)) \\ & + \beta(\psi(\omega_{\lambda}^G(T^n x_0, u, u)))\psi(\omega_{\lambda}^G(T^n x_0, TT^n x_0, TT^n x_0)) \\ & + \gamma(\psi(\omega_{\lambda}^G(T^n x_0, u, u)))\psi(\omega_{\lambda}^G(u, Tu, Tu)) \\ & + \psi(\omega_{\lambda}^G(T^n x_0, T^{n+1}x_0, T^{n+1}x_0)) \\ & = \alpha(\psi(\omega_{\lambda}^G(T^n x_0, u, u)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n x_0, u, u)) \\ & + \beta(\psi(\omega_{\lambda}^G(T^n x_0, u, u)))\psi(\omega_{\lambda}^G(T^n x_0, T^{n+1}x_0, T^{n+1}x_0)) \\ & + \gamma(\psi(\omega_{\lambda}^G(T^n x_0, u, u)))\psi(\omega_{\lambda}^G(u, Tu, Tu)) \\ & + \psi(\omega_{\lambda}^G(T^n x_0, T^{n+1}x_0, T^{n+1}x_0)), \end{aligned} \quad (3.23)$$

as $n \rightarrow \infty$, we have that

$$\psi(\omega_{\lambda}^G(u, Tu, Tu)) \leq \gamma(0)\psi(\omega_{\lambda}^G(u, Tu, Tu)) \quad (3.24)$$

for all $\lambda > 0$, which implies that

$$(1 - \gamma(0))\omega_{\lambda}^G(u, Tu, Tu) \leq 0 \quad \forall \lambda > 0. \quad (3.25)$$

Therefore,

$$\omega_{\lambda}^G(u, Tu, Tu) = 0 \quad \forall \lambda > 0, \quad (3.26)$$

where $1 - \gamma(0) < 1$. Hence, u is a fixed point of T for all $\lambda > 0$, i.e; $Tu = u$.

Finally, for the uniqueness, we can see from above that T has a fixed point $u \in X_{\omega^G}$. Suppose that there is another fixed point of T i.e; $Tv = v$, for $v \in X_{\omega^G}$, thus condition (2) of

Theorem 3.1 tells us that if $z \in X_{\omega^c}$ with $z \leq Tz$ and it is comparable to both u and v and $T^n z$ is also comparable to u and v for each $n \in \mathbb{N}$. Now for $\lambda > 0$, then $\psi(\omega_\lambda^G(T^{n+1}z, u, u))$ and $\psi(\omega_\lambda^G(T^{n+1}z, v, v))$ are finite. Claim : $u = v$. Indeed, using inequality (3.1), we have by taking $x = T^n z$ and $y = u$. First consider $\psi(\omega_\lambda^G(T^{n+1}z, u, u)) < \infty$, so that we have the following inequality by using condition 6 of **Proposition 2.2**

$$\begin{aligned}
 & \psi(\omega_\lambda^G(T^{n+1}z, u, u)) = \psi(\omega_\lambda(T^{n+1}z, Tu, Tu)) \\
 & = \psi(\omega_\lambda^G(TT^n z, Tu, Tu)) \\
 & \leq \alpha(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, u, u)) \\
 & + \beta(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(T^n z, TT^n z, TT^n z)) \\
 & + \gamma(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(u, Tu, Tu)) \\
 & = \alpha(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, u, u)) \\
 & + \beta(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(T^n z, T^{n+1}z, T^{n+1}z)) \\
 & \leq \alpha(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, u, u)) \\
 & + \psi\left(\omega_{\lambda/4}^G(T^{n+1}z, u, u)\right) + \psi\left(\omega_{\frac{\lambda}{4}}^G(T^{n+1}z, u, u)\right) \\
 & = \alpha(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, u, u)) \\
 & + \beta(\psi(\omega_\lambda^G(T^n z, u, u)))\left(\psi\left(\omega_{\lambda/2}^G(T^n z, u, u)\right)\right. \\
 & \left.+ 2\psi\left(\omega_{\lambda/4}^G(T^{n+1}z, u, u)\right)\right) \\
 & \leq \alpha(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(T^n z, u, u)) \\
 & + \beta(\psi(\omega_\lambda^G(T^n z, u, u)))\left(\psi(\omega_\lambda^G(T^n z, u, u))\right. \\
 & \left.+ 2\psi(\omega_\lambda^G(T^{n+1}z, u, u))\right) \\
 & = \alpha(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(T^n z, u, u)) \\
 & + \beta(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(T^n z, u, u)) \\
 & + 2\beta(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(T^{n+1}z, u, u)),
 \end{aligned} \tag{3.27}$$

so, we have that

$$\begin{aligned}
 \psi(\omega_\lambda^G(T^{n+1}z, u, u)) & \leq \frac{\alpha(\psi(\omega_\lambda^G(T^n z, u, u))) + \beta(\psi(\omega_\lambda^G(T^n z, u, u)))}{1 - 2\beta(\psi(\omega_\lambda^G(T^n z, u, u)))}\psi(\omega_\lambda^G(T^n z, u, u)) \\
 & \leq \psi(\omega_\lambda^G(T^n z, u, u)) \\
 & \leq \psi(\omega_\lambda^G(T^{n-1}z, u, u)) \\
 & \vdots \\
 & \leq \psi(\omega_\lambda^G(z, u, u)) \\
 & < \infty.
 \end{aligned} \tag{3.28}$$

Therefore, $\{\psi(\omega_\lambda^G(T^{n+1}z, u, u))\}_{n \geq 1}$ is nonincreasing sequence which is bounded below and converges to some real number $\ell \in [0, \infty)$. Assume that $\ell > 0$, using the fact that $\lim_{n \rightarrow \infty} \omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0) = 0$ for all $\lambda > 0$, from inequality 3, we have that

$$\begin{aligned} \psi(\omega_\lambda^G(T^{n+1}z, u, u)) &\leq \alpha(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, u, u)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(T^n z, T^{n+1}z, T^{n+1}z)) \\ &\leq \alpha(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(T^n z, u, u)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^n z, u, u)))\psi(\omega_\lambda^G(T^n z, T^{n+1}z, T^{n+1}z)). \end{aligned} \quad (3.29)$$

Using inequality 3 and letting $n \rightarrow \infty$, inequality 3 becomes

$$1 \leq \liminf_{n \rightarrow \infty} \alpha(\psi(\omega_\lambda^G(T^n z, u, u))). \quad (3.30)$$

Thus, by condition 4 of [Proposition 2.3](#) we have that

$$\lim_{n \rightarrow \infty} \omega_\lambda^G(T^n z, u, u) = 0 \quad (3.31)$$

for all $\lambda > 0$. Therefore, $T^n z \rightarrow u$ as $n \rightarrow \infty$.

Secondly consider $\psi(\omega_\lambda^G(T^{n+1}z, v, v)) < \infty$, from inequality [\(3.1\)](#), we have by taking $x = T^n z$ and $y = v$, so that we have the following inequality by using condition 6 of [Proposition 2.2](#)

$$\begin{aligned} \psi(\omega_\lambda^G(T^{n+1}z, v, v)) &= \psi(\omega_\lambda(T^{n+1}z, Tv, Tv)) \\ &= \psi(\omega_\lambda^G(TT^n z, Tv, Tv)) \\ &\leq \alpha(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, v, v)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_\lambda^G(T^n z, TT^n z, TT^n z)) \\ &\quad + \gamma(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_\lambda^G(v, Tv, Tv)) \\ &= \alpha(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, v, v)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_\lambda^G(T^n z, T^{n+1}z, T^{n+1}z)) \\ &\leq \alpha(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, v, v)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^n z, v, v)))\left(\psi(\omega_{\lambda/2}^G(T^n z, v, v))\right) \\ &\quad + \psi(\omega_{\lambda/4}^G(T^{n+1}z, v, v)) + \psi(\omega_{\lambda/4}^G(T^{n+1}z, v, v)) \\ &= \alpha(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, v, v)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^n z, v, v)))\left(\psi(\omega_{\lambda/2}^G(T^n z, v, v))\right) \\ &\quad + 2\psi(\omega_{\lambda/4}^G(T^{n+1}z, v, v)) \leq \alpha(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_\lambda^G(T^n z, v, v)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^n z, v, v)))\left(\psi(\omega_\lambda^G(T^n z, v, v))\right) \\ &\quad + 2\psi(\omega_\lambda^G(T^{n+1}z, v, v)) = \alpha(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_\lambda^G(T^n z, v, v)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_\lambda^G(T^n z, v, v)) \\ &\quad + 2\beta(\psi(\omega_\lambda^G(T^n z, v, v)))\psi(\omega_\lambda^G(T^{n+1}z, v, v)). \end{aligned} \quad (3.32)$$

Therefore, we have

$$\begin{aligned}
 \psi(\omega_{\lambda}^G(T^{n+1}z, v, v)) &\leq \frac{\alpha(\psi(\omega_{\lambda}^G(T^n z, v, v))) + \beta(\psi(\omega_{\lambda}^G(T^n z, v, v)))}{1 - 2\beta(\psi(\omega_{\lambda}^G(T^n z, v, v)))} \psi(\omega_{\lambda}^G(T^n z, v, v)) \\
 &\leq \psi(\omega_{\lambda}^G(T^n z, v, v)) \\
 &\leq \psi(\omega_{\lambda}^G(T^{n-1}z, v, v)) \\
 &\vdots \\
 &\leq \psi(\omega_{\lambda}^G(z, v, v)) \\
 &< \infty.
 \end{aligned} \tag{3.33}$$

Hence, $\{\psi(\omega_{\lambda}^G(T^{n+1}z, v, v))\}_{n \geq 1}$ is nonincreasing sequence which is bounded below and converges to some real number $\ell_0 \in [0, \infty)$. Suppose that $\ell_0 > 0$, using the fact that $\lim_{n \rightarrow \infty} \omega_{\lambda}^G(T^n x_0, T^{n+1}x_0, T^{n+1}x_0) = 0$ for all $\lambda > 0$. From inequality 3, we have that

$$\begin{aligned}
 \psi(\omega_{\lambda}^G(T^{n+1}z, v, v)) &\leq \alpha(\psi(\omega_{\lambda}^G(T^n z, v, v))) \psi(\omega_{\lambda+\nu(\lambda)}^G(T^n z, v, v)) \\
 &\quad + \beta(\psi(\omega_{\lambda}^G(T^n z, v, v))) \psi(\omega_{\lambda}^G(T^n z, T^{n+1}z, T^{n+1}z)) \\
 &\leq \alpha(\psi(\omega_{\lambda}^G(T^n z, v, v))) \psi(\omega_{\lambda}^G(T^n z, v, v)) \\
 &\quad + \beta(\psi(\omega_{\lambda}^G(T^n z, v, v))) \psi(\omega_{\lambda}^G(T^n z, T^{n+1}z, T^{n+1}z)).
 \end{aligned} \tag{3.34}$$

Using inequality 3 and letting $n \rightarrow \infty$, inequality 3 becomes

$$1 \leq \liminf_{n \rightarrow \infty} \alpha(\psi(\omega_{\lambda}^G(T^n z, v, v))). \tag{3.35}$$

Thus, by condition 4 of [Proposition 2.3](#) we have that

$$\lim_{n \rightarrow \infty} \omega_{\lambda}^G(T^n z, v, v) = 0 \tag{3.36}$$

for all $\lambda > 0$. Therefore, $T^n z \rightarrow v$ as $n \rightarrow \infty$.

Suppose, if possible, that $\lim_{n \rightarrow \infty} T^n z$ exists and not unique. Let $\lim_{n \rightarrow \infty} T^n z = u$ and $\lim_{n \rightarrow \infty} T^n z = v$ as we have seen above, where $u \neq v$. For each $\lambda > 0$, $u \neq v \Rightarrow \omega_{\lambda}^G(u, v, v) > 0$. If we take $\psi(\epsilon_1) = \frac{1}{3} \psi(\omega_{\lambda}^G(u, v, v)) > 0$, then for $\lambda > 0$, $\lim_{n \rightarrow \infty} T^n z = u \Rightarrow$ given $\epsilon_1 > 0$, $\exists m_1 \in \mathbb{N}$ such that $\psi(\omega_{\lambda/2}^G(u, T^n z, T^n z)) < \psi(\epsilon_1)$ for $n > m_1$. Again, $\lim_{n \rightarrow \infty} T^n z = v \Rightarrow$ given $\epsilon_1 > 0$, $\exists m_2 \in \mathbb{N}$ such that $\psi(\omega_{\lambda/4}^G(v, T^n z, T^n z)) < \psi(\epsilon_1)$ for $n > m_2$. Set $m = \max\{m_1, m_2\}$, then for $n \geq m$, by condition 6 of [Proposition 2.2](#), we have

$$\begin{aligned}
 \psi(\omega_{\lambda}^G(u, v, v)) &\leq \psi(\omega_{\lambda/2}^G(u, T^n z, T^n z)) \\
 &\quad + 2\omega_{\lambda/4}^G(v, T^n z, T^n z) \leq \psi(\omega_{\lambda/2}^G(u, T^n z, T^n z)) \\
 &\quad + 2\psi(\omega_{\lambda/4}^G(v, T^n z, T^n z)) < \psi(\epsilon_1) + 2\psi(\epsilon_1) \\
 &= 3\psi(\epsilon_1),
 \end{aligned}$$

which shows that $\psi(\omega_\lambda^G(u, v, v)) < \psi(\omega_\lambda^G(u, v, v))$ for all $\lambda > 0$. This is a contradiction. Hence, $u = v$. Therefore the fixed point of T is unique and the proof is complete. $\square$

We shall give an example to support Theorem 3.1 above.

Example 3.1. Let $X = \mathbb{R}$, define modular G -metric by $\omega_\lambda^G(x, y, y) = \infty$ if $\lambda \leq 2|x - y|$ and $\omega_\lambda^G(x, y, y) = 0$ if $\lambda > 2|x - y|$, and $\omega_\lambda^G(x, y, y) = \frac{G(x, y, y)}{\lambda}$, where $G(x, y, y) = 2|x - y|$ or $G(x, y, z) = |x - y| + |y - z| + |x - z|$ for $x, y, z \in \mathbb{R}$. We can see that $X_{\omega^G} = \mathbb{R}$. So it follows from Theorem 3.1 that $\mathbb{R}$ is a complete preordered modular G -metric space. Now define a map $T : \mathbb{R} \rightarrow \mathbb{R}$ by $Tx = \frac{x^3}{1+x^2}$. For $x, y \in \mathbb{R}$, then $\omega_\lambda^G(x, y, y) = \infty$ if $\lambda \leq 2|x - y|$, so inequality (3.1) is satisfied. Again, if $\lambda > 2|x - y|$ and $x, y \in \mathbb{R}$, then

$$G(Tx, Ty, Ty) = 2 \left| \frac{x^3}{1+x^2} - \frac{y^3}{1+y^2} \right| \leq 4|x - y| < 2\lambda, \text{ therefore, } \omega_\lambda^G(Tx, Ty, Ty) = 2\omega_\lambda^G(x, y, y) \leq 0.$$

We can take $\psi(t) = t$, $\alpha(t) = \beta(t) = \gamma(t) = \frac{1}{2}$. But T has a fixed point at $x = 0$.

Corollary 3.2. Let (X, ω^G) be a complete modular G -metric space with a preorder, $\leq$ and a nondecreasing self-mapping $T : X_{\omega^G} \rightarrow X_{\omega^G}$ on X_{ω^G} such that for each $\lambda > 0$, there is $\nu(\lambda) \in [0, \lambda)$ such that the following conditions hold:

(1)

$$\psi(\omega_\lambda^G(Tx, Ty, Ty)) \leq \alpha(\psi(\omega_\lambda^G(x, y, y)))\psi(\omega_{\lambda+\nu(\lambda)}^G(x, y, y)), \quad (3.37)$$

where $\psi \in \bar{\Psi}$ and $\alpha \in \mathcal{S}_{Ger}$ and distinct $x, y \in X_{\omega^G}$. Assuming that if a nondecreasing sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to x^* , then $x_n \leq x^*$ for each $n \in \mathbb{N}$,

(2) if ψ is subadditive and for any $x, y \in X_{\omega^G}$, there exists $z \in X_{\omega^G}$ with $z \leq Tz$ and $\omega_\lambda^G(z, Tz, Tz)$ is finite for all $\lambda > 0$ such that z is comparable to both x and y . Then T has a fixed point $u \in X_{\omega^G}$ and the sequence define by $\{T^n x_0\}_{n \geq 1}$ converges to u . Moreover, the fixed point of T is unique.

Proof. Let $x_0 \in X_{\omega^G}$ be such that $x_0 \leq Tx_0$ and let $x_n = Tx_{n-1} = T^n x_0$ for all $n \in \mathbb{N}$. Regarding that T is nondecreasing mapping, we have that $x_0 \leq Tx_0 = x_1$, which implies that $x_1 = Tx_0 \leq Tx_1 = x_2$. Inductively, we have

$$x_0 \leq x_1 \leq x_2 \leq \dots \leq x_{n-1} \leq x_n \leq x_{n+1} \leq \dots \quad (3.38)$$

Assume that there exists $n_0 \in \mathbb{N}$ such that $x_{n_0} = x_{n_0+1}$. Since $x_{n_0} = x_{n_0+1} = Tx_{n_0}$, then x_{n_0} is the fixed point of T . Now suppose that $x_n \not\leq x_{n+1}$ for all $n \in \mathbb{N}$, thus by inequality (3.38), we have that

$$x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n < x_{n+1} < \dots \quad (3.39)$$

Now for each $\lambda > 0$, and $x_0 < Tx_0$ for all $n \in \mathbb{N}$ implies that $\omega_\lambda^G(x_0, Tx_0, Tx_0) > 0$. Again, let $x_0 \in X_{\omega^G}$ such that $\omega_\lambda^G(x_0, Tx_0, Tx_0) < \infty \forall \lambda > 0$.

First, we show that for all $n \in \mathbb{N}$, the sequence $\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0) = 0$ for all $\lambda > 0$, as $n \rightarrow \infty$.

Assume that for each $n \in \mathbb{N}$, there exists $\lambda_n > 0$ such that $\omega_{\lambda_n}^G(T^{n+1} x_0, T^n x_0, T^n x_0) \neq 0$. Otherwise we are done. Suppose that for each $n \geq 1$, if $0 < \lambda < \lambda_n$, then we have $\omega_\lambda^G(T^{n+1} x_0, T^n x_0, T^n x_0) \neq 0$. Since $T^n x_0 \leq T^{n+1} x_0$, we have from inequality (3.37) that $\psi(\omega_{\lambda_n}^G(T^{n+1} x_0, T^n x_0, T^n x_0)) \leq \psi(\omega_\lambda^G(T^{n+1} x_0, T^n x_0, T^n x_0)) = \psi(\omega_\lambda^G(TT^n x_0, TT^{n-1} x_0, TT^{n-1} x_0))$. Take $x = T^n x_0$ and $y = T^{n-1} x_0$, then inequality (3.37) becomes;

$$\begin{aligned}
 & \psi(\omega_\lambda^G(T^{n+1}x_0, T^n x_0, T^n x_0)) \leq \alpha(\psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0))) \\
 & \times \psi(\omega_{\lambda+\nu(\lambda)}^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0)) \\
 & \leq \alpha(\psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0))) \\
 & \times \psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0)) \\
 & < \psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0)).
 \end{aligned} \tag{3.40}$$

Therefore, $\{\psi(\omega_\lambda^G(T^{n+1}x_0, T^n x_0, T^n x_0))\}_{n \geq 1}$ is nonincreasing and bounded below and converges to some real number $\tau \geq 0$. Assume that $\tau > 0$. In such a case,

$$\begin{aligned}
 \tau & < \psi(\omega_\lambda^G(T^{n+1}x_0, T^n x_0, T^n x_0)) \\
 & \leq \alpha(\psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0)))\psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0)) \\
 & < \psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0)),
 \end{aligned} \tag{3.41}$$

which implies that

$$\begin{aligned}
 1 & < \frac{\psi(\omega_\lambda^G(T^{n+1}x_0, T^n x_0, T^n x_0))}{\tau} \\
 & \leq \frac{\alpha(\psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0)))\psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0))}{\tau} \\
 & < \frac{\psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0))}{\tau}.
 \end{aligned} \tag{3.42}$$

Letting, $n \rightarrow \infty$, Theorem 2.1 ensure that $\{\alpha(\psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0)))\}_{n \geq 1} \rightarrow 1$ or from inequality 3,

$$1 \leq \liminf_{n \rightarrow \infty} \alpha(\psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0))). \tag{3.43}$$

As $\alpha \in \mathcal{S}_{Ger}$, then

$$\lim_{n \rightarrow \infty} \psi(\omega_\lambda^G(T^n x_0, T^{n-1}x_0, T^{n-1}x_0)) \rightarrow 0 \tag{3.44}$$

for all $\lambda > 0$, which contradicts the fact that $\tau > 0$. Thus $\tau = 0$, so that

$$\lim_{n \rightarrow \infty} \omega_\lambda^G(T^{n+1}x_0, T^n x_0, T^n x_0) = 0. \tag{3.45}$$

Hence $\omega_\lambda^G(T^{n+1}x_0, T^n x_0, T^n x_0) = 0$ for all $\lambda > 0, n \geq 1$. Following the proof of Theorem 3.1 above, we see that T has a unique fixed point in X_{ω^G} . $\square$

Theorem 3.3. Let (X, ω^G) be a complete modular G -metric space with a preorder, $\leq$, and a nondecreasing self-mapping $T : X_{\omega^G} \rightarrow X_{\omega^G}$ on X_{ω^G} such that for each $\lambda > 0$, there is $\nu(\lambda) \in [0, \lambda)$ such that the following conditions hold:

(1)

$$\psi(\omega_\lambda^G(T^m x, T^m y, T^m y)) \leq \alpha(\psi(\omega_\lambda^G(x, y, y)))\psi(\omega_{\lambda+\nu(\lambda)}^G(x, y, y)), \tag{3.46}$$

where $\psi \in \overline{\Psi}$ and $\alpha \in \mathcal{S}_{Ger}$ and distinct $x, y \in X_{\omega^G}$. Assuming that if a nondecreasing sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to x^* , then $x_n \leq x^*$ for each $n \in \mathbb{N}$,

(2) if ψ is subadditive and for any $x, y \in X_{\omega^G}$, there exists $z \in X_{\omega^G}$ with $z \leq Tz$ and $\omega_\lambda^G(z, Tz, Tz)$ is finite for all $\lambda > 0$ such that z is comparable to both x and y . Then T

has a fixed point $u \in X_{\omega^G}$ for some positive integer m and the sequence define by $\{T^n x_0\}_{n \geq 1}$ converges to u . Moreover, the fixed point of T is unique.

Proof. By Corollary 3.2, T^m has a fixed point say $u \in X_{\omega^G}$ for some positive integer $m \geq 1$. Now $T^m(Tu) = T^{m+1}u = T(T^m u) = Tu$, so Tu is a fixed point of T^m . By the uniqueness of fixed point of T^m , we have $Tu = u$. Therefore, u is a fixed point of T . Since fixed point of T is also fixed point of T^m , hence T has a unique fixed point in X_{ω^G} . $\square$

Theorem 3.4. Let (X, ω^G) be a complete modular G -metric space with a preorder, $\leq$ and a nondecreasing self-mapping $T: X_{\omega^G} \rightarrow X_{\omega^G}$ on X_{ω^G} such that for each $\lambda > 0$, there is $\nu(\lambda) \in [0, \lambda)$ such that the following conditions hold:

(1)

$$\begin{aligned} \psi(\omega_\lambda^G(T^m x, T^m y, T^m y)) &\leq \alpha(\psi(\omega_\lambda^G(x, y, y)))\psi(\omega_{\lambda+\nu(\lambda)}^G(x, y, y)) + \beta(\psi(\omega_\lambda^G(x, y, y))) \\ &\times \psi(\omega_\lambda^G(x, T^m x, T^m x)) + \gamma(\psi(\omega_\lambda^G(x, y, y)))\psi(\omega_\lambda^G(y, T^m y, T^m y)), \end{aligned} \quad (3.47)$$

where $\psi \in \bar{\Psi}$ and $\{\alpha, \beta, \gamma\} \in \mathcal{S}_{Ger}$ with $\alpha(t) + 2\max\{\sup_{t \geq 0} \beta(t), \sup_{t \geq 0} \gamma(t)\} < 1$, and distinct $x, y \in X_{\omega^G}$. Assuming that if a nondecreasing sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to x^* , then $x_n \leq x^*$ for each $n \in \mathbb{N}$,

(2) if ψ is subadditive and for any distinct $x, y \in X_{\omega^G}$, there exists $z \in X_{\omega^G}$ with $z \leq Tz$ and $\omega_\lambda^G(z, Tz, Tz)$ is finite for all $\lambda > 0$ such that z is comparable to both x and y . Then T has a fixed point $u \in X_{\omega^G}$ for some positive integer $m \geq 1$ and the sequence define by $\{T^n x_0\}_{n \geq 1}$ converges to u . Moreover, the fixed point of T is unique.

Proof. By Theorem 3.1, T^m has a fixed point say $u_* \in X_{\omega^G}$ for some positive integer $m \geq 1$. Now $T^m(Tu_*) = T^{m+1}u_* = T(T^m u_*) = Tu_*$, so Tu_* is a fixed point of T^m . By the uniqueness of fixed point of T^m , we have $Tu_* = u_*$. Therefore, u_* is a fixed point of T . Since fixed point of T is also fixed point of T^m , hence T has a unique fixed point in X_{ω^G} . $\square$

Theorem 3.5. Let (X, ω^G) be a complete modular G -metric space with a preorder, $\leq$ and a nondecreasing self-mapping $T: X_{\omega^G} \rightarrow X_{\omega^G}$ on X_{ω^G} such that for each $\lambda > 0$, there is $\nu(\lambda) \in [0, \lambda)$ such that the following conditions hold:

(1)

$$\begin{aligned} \psi(\omega_\lambda^G(Tx, Ty, Tz)) &\leq \alpha(\psi(\omega_\lambda^G(x, y, z)))\psi(\omega_{\lambda+\nu(\lambda)}^G(x, y, z)) + \beta(\psi(\omega_\lambda^G(x, y, z))) \\ &\times \psi(\omega_\lambda^G(x, Tx, Tx)) + \gamma(\psi(\omega_\lambda^G(x, y, z)))\psi(\omega_\lambda^G(y, Ty, Ty)) + \delta(\psi(\omega_\lambda^G(x, y, z))) \\ &\times \psi(\omega_\lambda^G(z, Tz, Tz)), \end{aligned} \quad (3.48)$$

where $\psi \in \bar{\Psi}$ and $\{\alpha, \beta, \gamma, \delta\} \in \mathcal{S}_{Ger}$ with $\alpha(t) + 2\max\{\sup_{t \geq 0} \beta(t), \sup_{t \geq 0} \gamma(t), \sup_{t \geq 0} \delta(t)\} < 1$, and $x, y, z \in X_{\omega^G}$. Assuming that if a nondecreasing sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to x , then $x_n \leq x$ for each $n \in \mathbb{N}$,

(2) if ψ is subadditive and for any $x, y, z \in X_{\omega^G}$, there exists $w \in X_{\omega^G}$ with $w \leq Tw$ and $\omega_\lambda^G(w, Tw, Tw)$ is finite for all $\lambda > 0$ such that w is comparable to both x, y and z . Then T has a fixed point $u \in X_{\omega^G}$ and the sequence define by $\{T^n x_0\}_{n \geq 1}$ converges to u . Moreover, the fixed point of T is unique.

Proof. Let $x_0 \in X_{\omega^G}$ be such that $x_0 \leq Tx_0$ and let $x_n = Tx_{n-1} = T^n x_0$ for all $n \in \mathbb{N}$. Regarding that T is nondecreasing mapping, we have that $x_0 \leq Tx_0 = x_1$, which implies that $x_1 = Tx_0 \leq Tx_1 = x_2$. Inductively, we have

$$x_0 \leq x_1 \leq x_2 \leq \cdots \leq x_{n-1} \leq x_n \leq x_{n+1} \leq \cdots \quad (3.49)$$

Assume that there exists $n_0 \in \mathbb{N}$ such that $x_{n_0} = x_{n_0+1}$. Since $x_{n_0} = x_{n_0+1} = Tx_{n_0}$, then x_{n_0} is the fixed point of T . Now suppose that $x_n \not\leq x_{n+1}$ for all $n \in \mathbb{N}$, thus by inequality (3.38), we have that

$$x_0 < x_1 < x_2 < \cdots < x_{n-1} < x_n < x_{n+1} < \cdots \quad (3.50)$$

Now for each $\lambda > 0$, and $x_0 < Tx_0$ for all $n \in \mathbb{N}$ implies that $\omega_\lambda^G(x_0, Tx_0, Tx_0) > 0$. Again, let $x_0 \in X_{\omega^G}$ such that $\omega_\lambda^G(x_0, Tx_0, Tx_0) < \infty \forall \lambda > 0$.

First, we show that for all $n \in \mathbb{N}$, the sequence $\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0) = 0$ for all $\lambda > 0$, as $n \rightarrow \infty$.

Assume that, for each $n \in \mathbb{N}$, there exists $\lambda_n > 0$ such that $\omega_{\lambda_n}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0) \neq 0$. Otherwise there is nothing to prove. Suppose that for each $n \geq 1$, if $0 < \lambda < \lambda_n$, then we have that $\omega_\lambda^G(T^n x_0, T^n x_0, T^{n+1} x_0) \neq 0$. Since $T^n x_0 \leq T^{n+1} x_0$, we have from inequality (3.5) that $\psi(\omega_{\lambda_n}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)) \leq \psi(\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)) = \psi(\omega_\lambda^G(TT^{n-1} x_0, TT^n x_0, TT^n x_0))$. Take $x = T^{n-1} x_0$ and $y = T^n x_0 = z$, then inequality (3.5) becomes;

$$\begin{aligned} \psi(\omega_{\lambda_n}^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)) &\leq \psi(\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)) \\ &\leq \alpha(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_{\lambda+\nu(\lambda)}^G(T^{n-1} x_0, T^n x_0, T^n x_0)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^{n-1} x_0, TT^{n-1} x_0, TT^{n-1} x_0)) \\ &\quad + \gamma(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^n x_0, TT^n x_0, TT^n x_0)) \\ &\quad + \delta(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^n x_0, TT^n x_0, TT^n x_0)) \\ &= \alpha(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_{\lambda+\nu(\lambda)}^G(T^{n-1} x_0, T^n x_0, T^n x_0)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0)) \\ &\quad + \gamma(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)) \\ &\quad + \delta(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)) \\ &\leq \alpha(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0)) \\ &\quad + \beta(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0)) \\ &\quad + \gamma(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)) \\ &\quad + \delta(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) \times \psi(\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)), \end{aligned} \quad (3.51)$$

which implies that

$$\begin{aligned} \psi(\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0)) &\leq \rho \psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0)) \\ &\leq \psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0)) \\ &\quad \vdots \\ &\leq \psi(\omega_\lambda^G(x_0, Tx_0, Tx_0)) < \infty, \end{aligned} \quad (3.52)$$

where

$$\rho := \frac{\alpha(\psi(\omega_\lambda^G(T^{n-1}x_0, T^n x_0, T^n x_0))) + \beta(\psi(\omega_\lambda^G(T^{n-1}x_0, T^n x_0, T^n x_0)))}{1 - (\gamma(\psi(\omega_\lambda^G(T^{n-1}x_0, T^n x_0, T^n x_0))) + \delta(\psi(\omega_\lambda^G(T^{n-1}x_0, T^n x_0, T^n x_0))))}. \quad (3.53)$$

230

Therefore, $\{\psi(\omega_\lambda^G(T^n x_0, T^{n+1} x_0, T^{n+1} x_0))\}_{n \geq 1}$ is nonincreasing and bounded below, hence converges to some real number $s \geq 0$. We can also see clearly that by taking $\theta(\cdot) := \gamma(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0))) + \delta(\psi(\omega_\lambda^G(T^{n-1} x_0, T^n x_0, T^n x_0)))$, as $y = T^n x_0 = z$, then following [Theorem 3.1](#), T has a unique fixed point in X_{ω^G} . This complete the proof. $\square$

Theorem 3.6. Let (X, ω^G) be a complete modular G -metric space with a preorder, $\leq$ and a nondecreasing self-mapping $T : X_{\omega^G} \rightarrow X_{\omega^G}$ on X_{ω^G} such that for each $\lambda > 0$, there is $\nu(\lambda) \in [0, \lambda)$ such that the following conditions hold:

(1)

$$\begin{aligned} \psi(\omega_\lambda^G(T^m x, T^m y, T^m z)) &\leq \alpha(\psi(\omega_\lambda^G(x, y, z)))\psi(\omega_{\lambda+\nu(\lambda)}^G(x, y, z)) \\ &+ \beta(\psi(\omega_\lambda^G(x, y, z)))\psi(\omega_\lambda^G(x, T^m x, T^m x)) + \gamma(\psi(\omega_\lambda^G(x, y, z)))\psi(\omega_\lambda^G(y, T^m y, T^m y)) \\ &+ \delta(\psi(\omega_\lambda^G(x, y, z)))\psi(\omega_\lambda^G(z, T^m z, T^m z)), \end{aligned} \quad (3.54)$$

where $\psi \in \overline{\Psi}$ and $\{\alpha, \beta, \gamma, \delta\} \in \mathcal{S}_{Ger}$ with $\alpha(t) + 2\max\{\sup_{t \geq 0} \beta(t), \sup_{t \geq 0} \gamma(t), \sup_{t \geq 0} \delta(t)\} < 1$, and $x, y, z \in X_{\omega^G}$. Assuming that if a nondecreasing sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to x , then $x_n \leq x$ for each $n \in \mathbb{N}$;

(2) if ψ is subadditive and for any $x, y, z \in X_{\omega^G}$, there exists $w \in X_{\omega^G}$ with $w \leq Tw$ and $\omega_\lambda^G(w, Tw, Tw)$ is finite for all $\lambda > 0$ such that w is comparable to both x, y and z . Then T has a fixed point $u \in X_{\omega^G}$ for some positive integer $m \geq 1$ and the sequence define by $\{T^m x_0\}_{n \geq 1}$ converges to u . Moreover, the fixed point of T is unique.

Proof. Take $y = z$ and $\phi(\cdot) = \gamma(\psi(\omega_\lambda^G(x, y, z))) + \delta(\psi(\omega_\lambda^G(x, y, z)))$, then [Theorem 3.5](#) tells us that T^m has a fixed point say $u \in X_{\omega^G}$ for some positive integer $m \geq 1$. Therefore, [Theorem 3.4](#) shows that T has a unique fixed point in X_{ω^G} . $\square$

4. Applications to nonlinear Volterra-Fredholm-type integral equations

In this section, we construct a system of nonlinear integral equation that satisfies the conditions of [Theorem 3.1](#). We consider the following general nonlinear Volterra-Fredholm-type integral equations.

$$u(t, x) = h(t, x) + \int_0^t \int_B F(t, x, s, y, u(s, y), (L^* u)(s, y)) dy ds, \quad (4.1)$$

and

$$v(t, x) = e(t, x) + \int_0^t \int_B G(t, x, s, y, v(s, y), (L^* v)(s, y)) ds dy, \quad (4.2)$$

where

$$(L^*u)(t, x) = \int_0^t \int_B K(t, x, \tau, z, u(\tau, z)) dz d\tau, \quad (4.3)$$

and

$$(L^*v)(t, x) = \int_0^t \int_B K(t, x, \tau, z, v(\tau, z)) dz d\tau, \quad (4.4)$$

h, F, K and e, G, L are given functions and u, v are the unknown functions. We assume that $h, e \in C(\mathbb{R}_+ \times B, \mathbb{R}^n)$, $K \in C(\Omega \times \mathbb{R}^n, \mathbb{R}^n)$, $F \in C(\Omega \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n)$, $G \in C(\Omega \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n)$ and $\Omega = \{(t, x, s, y) : 0 \leq s \leq t < \infty; x, y \in B\}$, $B = \prod_{j=1}^n [a_j, b_j]$, $b_j > a_j$. Take $\sup_{(t,x)} \in \mathbb{R}_+ \times B g(s, y) \leq \frac{1}{t \prod_{j=1}^n [a_j, b_j]}$, where $(t, x) \in \mathbb{R}_+ \times B$.

Let $\alpha, \beta, \gamma > 0$ with $\alpha(t) + 2\max_{t \in \Omega} \{\sup_{t \geq 0} \beta(t), \sup_{t \geq 0} \gamma(t)\} < 1$ such that

$$\begin{aligned} & \|F(t, x, s, y, u(s, y), (L^*u)(s, y)) - G(t, x, s, y, v(s, y), (L^*v)(s, y))\| \\ & \leq g(s, y) \{ \alpha(\|u - v\|) \|u - v\| + \beta(\|u - v\|) m(u, L^*u) + \gamma(\|u - v\|) r(v, L^*v) \} \end{aligned}$$

Let $F, G : C(\Omega \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n) \rightarrow \mathbb{R}^n$ be such that $F_u, G_v \in C(\Omega \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n)$ and let

$$F_u = \int_0^t \int_B F(t, x, s, y, u(s, y), (L^*u)(s, y)) dy ds, \quad (4.5)$$

and

$$G_v = \int_0^t \int_B G(t, x, s, y, v(s, y), (L^*v)(s, y)) ds dy, \quad (4.6)$$

for $F \in C(\Omega \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n)$, $G \in C(\Omega \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n)$ and u, v are the unknown functions.

Now for any $\lambda > 0$, we define

$$\omega_\lambda^G(x, y, z) := \frac{1}{2(1 + \lambda)} \sup_{(t,x) \in \mathbb{R}_+ \times B} \{ \|x(t) - y(t)\| + \|y(t) - z(t)\| + \|x(t) - z(t)\| \}, \quad (4.7)$$

so that

$$\omega_\lambda^G(x, y, y) := \frac{1}{(1 + \lambda)} \sup_{(t,x) \in \mathbb{R}_+ \times B} \{ \|x(t) - y(t)\| \}. \quad (4.8)$$

In fact Eqns. (4.7) and (4.8) satisfies all the conditions in Definition 2.6 endowed with $X_{\omega^G} = (X, \omega^G) = C(\Omega \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n)$

Now, take $A = \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \{ \|F_u - G_v + h(t, x) - e(t, x)\| \}$, so that

$$\begin{aligned}
 A &\leq \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \left\{ \int_0^t \int_B \|F(t, x, s, y, u(s, y), (L^*u)(s, y)) \right. \\
 &\quad \left. - G(t, x, s, y, v(s, y), (L^*v)(s, y))\| \times dy ds \right\} \\
 &\leq \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \left\{ \int_0^t \int_B g(s, y) \{ \alpha(\|u - v\|) \|u - v\| \right. \\
 &\quad \left. + \beta(\|u - v\|) m(u, L^*u) + \gamma(\|u - v\|) r(v, L^*v) \} dy ds \right\} \\
 &\leq \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \{ \{ \alpha(\|u - v\|) \|u - v\| + \beta(\|u - v\|) m(u, L^*u) \\
 &\quad + \gamma(\|u - v\|) r(v, L^*v) \} \} \\
 &\leq \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \{ \alpha(\|u - v\|) \|u - v\| \} \\
 &\quad + \frac{1}{1+\lambda} \sup_{t \in \Omega} \{ \beta(\|u - v\|) m(u, L^*u) + \gamma(\|u - v\|) r(v, L^*v) \} \\
 &\leq \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \{ \alpha(\|u - v\|) \|u - v\| \} \\
 &\quad + \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \{ \beta(\|u - v\|) m(u, L^*u) \} \\
 &\quad + \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \{ \gamma(\|u - v\|) r(v, L^*v) \},
 \end{aligned} \tag{4.9}$$

where $m, r \in C(\mathbb{R}_+ \times B \times \mathbb{R}^n, \mathbb{R}^n)$

$$m(u, L^*u)(t, x) = \sup_{(t,x) \in \mathbb{R}_+ \times B} \|F_u(t, x) + h(t, x) - u(t, x)\|, \tag{4.10}$$

$$r(v, L^*v)(t, x) = \sup_{(t,x) \in \mathbb{R}_+ \times B} \|G_v(t, x) + e(t, x) - v(t, x)\| \tag{4.11}$$

Theorem 4.1. Let $X_{\omega^G} = C(\Omega \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n)$ be a complete modular G -metric space and $\omega^G : (0, \infty) \times X_{\omega^G} \times X_{\omega^G} \times X_{\omega^G} \rightarrow \mathbb{R}_+^n \cup \{\infty\}$ be defined by

$$\omega_\lambda^G(u, v, v) := \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \|u(t, x) - v(t, x)\|, \lambda > 0 \tag{4.12}$$

and $u \leq v \Leftrightarrow u(t, x) \leq v(t, x) \quad \forall (t, x) \in \mathbb{R}_+ \times B$. Let $F_u, G_v : C(\Omega \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n) \rightarrow \mathbb{R}^n$ are such that $F_u, G_v \in X_{\omega^G}$ for each $u, v \in X_{\omega^G}$ and F_u, G_v satisfying Eqns. (4.5) and (4.6), respectively, for all $(t, x) \in \mathbb{R}_+ \times B$. Suppose that there exists nonnegative reals $\alpha, \beta, \gamma > 0$ with $\alpha(t) + 2\max_{t \in \Omega} \{ \sup_{t \geq 0} \beta(t), \sup_{t \geq 0} \gamma(t) \} < 1$ such that inequality 4 is satisfied for every $u, v \in X_{\omega^G}$. Moreover if ψ is subadditive and for any $u, v \in X_{\omega^G}$, there exists $w_0, w_1 \in X_{\omega^G}$ with $w_0 \leq w_1$ and $\omega_\lambda^G(w_0, w_1, w_1)$ is finite for all $\lambda > 0$ such that w is comparable to both u and v . Then the system of integral Eqns (4.1) and (4.2) have a unique solution in X_{ω^G} .

Proof. Define the mapping $T : X_{\omega^G} \rightarrow X_{\omega^G}$ by $Tu = F_u + e$ and $Tv = G_v + h$. Then for $\lambda > 0$, $\omega_\lambda^G(Tu, Tv, Tv) = \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \backslash norm F_u(t, x) - G_v(t, x) + e(t, x) - h(t, x)$, $\omega_\lambda^G(u, Tu, Tu) = \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \backslash norm F_u(t, x) + e(t, x) - u(t, x)$ and $\omega_\lambda^G(v, Tv, Tv) = \frac{1}{1+\lambda} \sup_{(t,x) \in \mathbb{R}_+ \times B} \backslash norm G_v(t, x) + h(t, x) - v(t, x)$. So from inequality 4, we get by noticing that ψ is continuous and subadditive and there exists $\nu \in [0, \lambda)$ such that

$$\begin{aligned} \psi(\omega_\lambda^G(Tu, Tv, Tv)) &\leq \alpha(\psi(\omega_\lambda^G(u, v, v)))\psi(\omega_{\lambda+\nu(\lambda)}^G(u, v, v)) \\ &+ \beta(\psi(\omega_\lambda^G(u, v, v)))\psi(\omega_\lambda^G(u, Tu, Tu)) + \gamma(\psi(\omega_\lambda^G(u, v, v)))\psi(\omega_\lambda^G(v, Tv, Tv)), \end{aligned} \quad (4.13)$$

where $\psi \in \bar{\Psi}$. By Theorem 3.1, we conclude that the system of nonlinear Volterra-Fredholm integral Eqns (4.1) and (4.2) have a unique solution in X_{ω^G} . $\square$

References

- Geraghty M. On contractive mapping. Proc Amer Math Soc. 1973; 40: 604-8.
- Gordji ME, Cho YJ, Pirbavafa S. A generalization of Geraghty's theorem in partially ordered metric spaces and application to ordinary differential equations. Fixed Point Theor Appl. 2012; 74: 1687-812.
- Bhaskar IG, Lakshmikantham V. Fixed point theorems in partially ordered metric spaces and applications. Nonlinear Anal. 2006; 65: 1379-93.
- Yolacan E. Fixed point theorems for Geraghty type contraction mappings and coupled fixed point results in 0-complete ordered partial metric spaces. Int J Math Sci. 2016; 2016: 5.
- Faraji H, Savic D, Radenovic S. Fixed point theorems for Geraghty contraction type mappings in b-metric spaces and application. Axioms. 2019; 8: 34. doi: [10.3390/axioms/8010034](https://doi.org/10.3390/axioms/8010034).
- Gupta V, Shatanawi W, Mani N. Fixed point theorems for (ψ, β) -Geraghty contraction-type maps in ordered metric spaces and some applications to integral and ordinary differential equations. J Fixed Point Theor Appl. 2017; 19: 1251-1267.
- Chaipunya P, Cho YJ, Kumam P. Geraghty-type theorems in modular metric spaces with application to partial differential equation. Adv Differ Equ. 2012; 83: 1687-847.
- Agarwal RP, Karapinar E, O'Regan D, Roldan-Lopez-de-Hierro AF. Fixed point theory in metric type spaces, New York Dordrecht London: Springer Cham Heidelberg. 2015.
- Abbas M, Nazir T, Shatanawi W, Mustafa Z. Fixed and related fixed point theorems for three maps in G-metric spaces. Hacet J Math Stat. 2012; 41(2): 291-06.
- Chistyakov VV. Metric modular spaces, I basic concepts. Nonlinear Anal Theor Meth Appl. 2010; 72: 1-14.
- Chistyakov VV. Metric modular spaces, II Applications to superposition operators. Nonlinear Anal Theor Meth Appl. 2010; 72: 15-30.
- Nakano H. Modulare semi-ordered linear spaces. Vol 1, Tokyo: Maruzen. 1950.
- Orlicz W. Collected papers. Vols I, II, Warszawa: PWN. 1988.
- Musielak J. Orlicz spaces and modular spaces, Lecture Notes in Math. Vol. 1034, Berlin: Springer-verlag. 1983.
- Nakano H. Topology of linear topological spaces. Vol 3, Tokyo: Maruzen. 1951.
- Chistyakov VV. A fixed point theorem for contractions in metric modular spaces. arXiv:1112.5561. 2011; 65-92.
- Mutlu A, Ozkan K, Gurdal U. A new fixed point theorem in modular metric spaces. Int J Anal Appl. 2018; 4: 472-83.

18. Okeke GA, Kim JK. Approximation of common fixed point of three multi-valued ρ -quasi-nonexpansive mappings in modular function spaces. *Nonlinear Func Anal Appl.* 2019; 24(4): 651-64.
19. Rhoades BE. Two fixed point theorems for mappings satisfying a general contractive condition in integral-type. *Int J Math Sci.* 2003; 63: 4007-13.
20. Okeke GA, Khan SH. Approximation of fixed point of multivalued ρ -quasi-contractive mappings in modular function spaces. *Arab J Math Sci.* 2020; 26(1/2): 75-93.
21. Bishop SA, Okeke GA, Eke K. Mild solutions of evolution quantum stochastic differential equations with nonlocal conditions. *Math Meth Appl Sci.* 2020: 1-8. doi: [10.1002/mma.6368](https://doi.org/10.1002/mma.6368).
22. Chlebowicz A. Solvability of an infinite system of nonlinear integral equations of Volterra-Hammerstein type. *Adv. Nonlinear Anal.* 2020 9(1): 1187-204.
23. Liu Z, Li X, Kang SM, Cho YJ. Fixed point theorems for mappings satisfying contractive conditions of integral-type and applications. *Fixed Point Theor Appl.* 2011; 64: 1687-812.
24. Okeke GA, Francis D, de la Sen M. Some fixed point theorems for mappings satisfying rational inequality in modular metric spaces with applications. *Heliyon* 2020; 6: e04785.
25. Papageorgiou NS, Radulescu VR, Repovs DD. *Nonlinear analysis - theory and methods*, Cham: Springer Monographs in Mathematics. Springer. 2019.
26. Zhao HY. Pseudo almost periodic solutions for a class of differential equation with delays depending on state. *Adv Nonlinear Anal.* 2020; 9(1): 1251-58.
27. Azadifar B., Maramaei M., Sadeghi G.. On the modular G-metric spaces and fixed point theorems. *J Nonlinear Sci Appl.* 2013; 6: 293-304.
28. Bali NP. *Golden real analysis*, New Delhi: Laxmi Publications. 1986.
29. Schechter E. *Handbook of analysis and its foundations*. New York: Academic Press. 1997.

Further reading

30. Amini-Harandi A. A fixed point theory for generalized quasicontraction maps in vector modular spaces. *Comput Math Appl.* 2011; 61: 1891-97.
31. Mustafa Z, Khandagji M, Shatanawi W. Fixed point results on complete G-metric spaces. *Stud Sci Math Hung.* 2011: 48(3); 304-19.
32. Nakano H. On the stability of functional equations. *Aequationes Math.* 2009; 77: 33-88.

Corresponding author

Godwin Amechi Okeke can be contacted at: gaokeke1@yahoo.co.uk

For instructions on how to order reprints of this article, please visit our website:

www.emeraldgroupublishing.com/licensing/reprints.htm

Or contact us for further details: permissions@emeraldinsight.com