

Chapter 12

Primary Care for Adolescents

Pierre-André Michaud, Johanna P.M. Vervoort and Danielle Jansen

Abstract

Adolescence is a time when a young person develops his or her identity, acquires greater autonomy and independence, experiments and takes risks and grows mentally and physically. To successfully navigate these changes, an accessible and health system when needed is essential.

We assessed the structure and content of national primary care services against these standards in the field of adolescent health services. The main criteria identified by adolescents as important for primary care are as follows: accessibility, staff attitude, communication in all its forms, staff competency and skills, confidential and continuous care, age appropriate environment, involvement in health care, equity and respect and a strong link with the community.

We found that although half of the Models of Child Health Appraised countries have adopted adolescent-specific policies or guidelines, many countries do not meet the current standards of quality health care for adolescents. For example, the ability to provide emergency mental health care or respond to life-threatening behaviour is limited. Many countries provide good access to contraception, but specialised care for a pregnant adolescent may be hard to find.

Access needs to be improved for vulnerable adolescents; greater advocacy should be given to adolescent health and the promotion of good health habits. Adolescent health services should be well publicised, and adolescents need to feel empowered to access them.

Keywords: Adolescents; health care; preventive care; primary care services; mental health; sexual and reproductive health



© European Commission. Published by Emerald Publishing Limited. This chapter is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this chapter (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

Introduction

Adolescents (defined in this survey as individuals aged 10–19 years) have specific needs compared with younger children. They are in the process of developing their identity and acquiring autonomy, their bodies and minds are growing, and it is a time of experimenting and risk-taking, and increasing independence (Jansen et al., 2018; Michaud, Blum, & Ferron, 1998; WHO, 2014b). Adolescents need to feel confident in their ability to access primary care services, in the form of advice, prevention and treatment services – independently of their parents or guardians if appropriate (Michaud et al., 2010). Models of Child Health Appraised (MOCHA) has identified young people as an important group in terms of their health and also in terms of children’s rights (United Nations General Assembly, 1990). Adolescents should be respected and involved as much as possible in all decisions regarding their life and their health. To provide optimal services, the primary healthcare system and the health professionals providing services need to recognise the needs of adolescents and adapt policies accordingly (Sawyer et al., 2014).

The health of an adolescent depends on many factors that lie beyond the healthcare system, such as the economic situation of the country, the climate and the culture, the organisation of the educational system, the presence or absence of preventive activities and so on (see Chapter 17; Patton et al., 2012; Patton et al., 2016; World Health Organization, 2014b; World Health Organization, 2017; WHO, 2014a). MOCHA investigated the extent to which the current health systems of European countries met the healthcare needs of adolescents aged 10–18, as being the upper age of childhood as defined by the Convention on the Rights of the Child (CRC) (United Nations, 1989).

There are models of quality health care available for adolescents (Michaud & Baltag, 2015; Michaud, Weber, Namazova-Baranova, & Ambresin, 2018; World Health Organization, 2014; World Health Organization, 2015a; World Health Organization, 2015b; World Health Organization, 2015c), most of which refer to the concept of adolescent/youth-friendly health services and care jointly developed by the World Health Organization (WHO), United Nations Children’s Fund (UNICEF) and United Nations Population Fund (UNFPA). These models have also been validated by young people themselves (Ambresin, Bennett, Patton, Sancu, & Sawyer, 2013), as a result of surveys about the main ingredients of fair and high-quality health services and care. The main criteria mentioned by young people in this survey are as follows:

- accessibility (flexible schedule, possibility to drop in), location (public transportation), affordability (financial coverage) and equity;
- staff attitude: respectful, supportive, empathetic, trustworthy and honest;
- communication: developmentally appropriate, understandable, active listening and provision of information;
- staff competency and skills, both technical and medical (health care), and comprehensive and holistic approach (multi professional: e.g. providing

curative and preventive services in the broad area of adolescent health, including mental health, substance use, sexual & reproductive health) (see Chapter 13);

- guideline-driven care: confidentiality, autonomy, privacy and continuity of care;
- age appropriate environment: clean and teen-oriented physical space, health information, access to internet, pamphlets and leaflets;
- involvement in health care, participation, shared decision-making approach and continuity of care;
- equity and respect of adolescents' rights (CRC); and
- link with the community, networking approach and community support;

These comments align closely with MOCHA findings from young people about their experiences of primary care (see Chapter 3), and MOCHA sought to address whether the experience of primary healthcare services met these standards.

This report complements the survey on school health services, as described in Chapter 11, and assesses the extent to which the structure and content of primary care services comply with available standards in the field of adolescent health care.

Methods

We created a questionnaire on adolescent primary care services to be sent to the MOCHA country agents (see Chapter 1). The questionnaire was divided into three sections and contained 43 questions on structural and content issues that are specific to adolescent care. Each section began with a typical clinical vignette to assist the Country Agent in understanding what information was expected. These included the existence of guidelines or policies regarding adolescent-friendly health services and care, the respect of adolescent rights, access of adolescents to appropriate health care as well as the continuity of care. The two last sections of the questionnaire focussed on two major healthcare areas during adolescence: mental health and self-harm, and sexual and reproductive health. Complete data from all thirty countries were available for analysis.

Results

Adolescent Primary Care Services

We assessed the country agent answers against the existing adolescent-friendly health services and care (AFHSC) guidelines. Thirteen out of the 30 countries surveyed indicated that they were aware of and follow the AFHSC guidelines, and a document to this end is available nationally. However, it was impossible to ascertain whether the documents are applied and to what extent. One of the questions tackled the existence of specialised services for adolescents. More than half the countries (16/30) have set up such specialised centres to deliver adolescent health care, although these are likely to be in selected cities and not in all regions of a country. Some units address specific issues (such as sexual




Figure 12.1. Countries with extensive policy on AHS.

and reproductive life or mental health), and some are more broadly oriented. Many, if not most, are run by multidisciplinary team ($N = 16$), and in eleven countries, the country agents claim that professionals in charge have received formal training in adolescent health (see Chapter 13). Figure 12.1 shows the countries that have an extensive policy on adolescent health services as recommended currently (Kokotailo et al., 2018). For more information, see Jansen et al. (2018).

Adolescents' Rights and Ethical Issues

The respect of confidentiality and privacy is, according to young people, of utmost importance, and this applies to all countries of the world (see Chapter 3; Baltag & Mathison, 2010; Bell, Breland, & Ott, 2013; Committee on Adolescence, 2016; National Research Council and Institute of Medicine, 2009; World Health Organization, 2016; Michaud et al., 2010). Indeed, when it comes to discussing sensitive issues such as sexual conduct or contraception, risk-taking, problematic eating patterns or substance use, young people need to be confident that the healthcare professional will not disclose information unless the situation is life threatening or unless the adolescent feels comfortable to disclose. However, the right to confidentiality is linked with the young person's decision-making capacity (competence), and healthcare providers are not necessarily well equipped to assess such a capacity (Michaud, Blum, Benaroyo,

Zermatten, & Baltag, 2015). In 13 out of the thirty countries surveyed, the existence of a formal legislation or policy tackling the issue of confidentiality was confirmed by country agents. Five countries also have policies but restricted by an age range. In only nine countries, guidelines are available about how to assess a young person's competence. Another important aspect of confidentiality is the right of a young person to access health care without the knowledge of parents. In 20 countries, adolescents have the right to consult a doctor without parents (or any substitute) knowing, and in around the same proportion of countries, adolescents have the right to choose their doctor themselves ($N = 17$). Finally, shared decision-making, for example the right to refuse a treatment or choosing another alternative than the one chosen by the parents, is a right that should be given to competent young patients. In around half of the countries ($N = 9$), a policy exists to guarantee such a right.

Access to Health Care

Access to health care is an important issue for adolescents. Blair, Rigby, and Alexander (2017) stated that most European countries provide some kind of sustainable insurance system that covers the healthcare expenditures of children and young people. The potentially limiting factors to access of adolescents to health care is thus more likely linked to a lack of knowledge of what exists and where to be able to consult freely and expect high-quality health care. In addition to this, it is sometimes difficult to access services because of a lack of availability, due to under-resourcing or a shortage of health professionals skilled in adolescent health care. This is particularly pertinent to so-called vulnerable adolescents, such as migrants and adolescents from deprived socio-economic background or 'drop-out' adolescents who are homeless or in temporary accommodation. Globally, around 50% of countries have developed policies or strategies that aim to improve access to care for adolescents facing situations of vulnerability. In around half of the countries ($N = 16$), it is possible for adolescents in such situations to consult primary care spontaneously. Half of the countries are able to offer translators if needed, at least in some regions, and/or to provide professionals who have an expertise in cross-cultural issues. Moreover, just about two thirds of the countries ($N = 20$) have policies which encourage an inter-professional approach to disruptive behaviour adolescents having left or being about to leave the mainstream educational system (see Chapter 11).

Access to Mental Health Care and Sexual and Reproductive Health Care

Issues such as conduct disorders, violence, depression and self-harm/suicide are increasingly recognised as important threats to adolescents' mental health (Nair et al., 2015; Patton et al., 2012; Potrebny, Wiium, & Lundegard, 2017; Steinberg et al., 2017). The majority of countries ($N = 25$) have some kind of suicide prevention programme, and a similar number are able to provide same-day referral appointments for suicide or mental health breakdown, but only half ($N = 14$) of the surveyed countries provide guidelines to primary care physicians as how to screen for mental health problems and disorders in adolescents, and only

Table 12.1. Indicators of quality management for mental health services and sexual and reproductive health care for adolescents.

Country	Quality Management Infrastructure	
	Guidelines PC Screening Young People on Mental Health Issues	Guidelines/Standards for PC Professionals about Adolescent Pregnancy
Austria	No	No
Belgium-F	Unclear	No
Bulgaria	Yes	No
Croatia	Yes	No
Cyprus	No	No
Czech Republic	Yes	No
Denmark	No	Yes
Estonia	Yes	Yes
Finland	Yes	Yes
France	Unclear	Yes
Germany	Yes	No
Greece	Yes	No
Hungary	No	—
Iceland	No	No
Ireland	Yes	Yes
Italy	Yes	No
Latvia	No	No
Lithuania	Unclear	No
Luxembourg	No	No
Malta	No	No
Netherlands	Yes	No
Norway	No	No
Poland	No	Yes
Portugal	Yes	No
Romania	No	No
Slovakia	No	No
Slovenia	Yes	No
Spain	Yes	Yes
Sweden	No	No
UK ENG	Yes	Yes

fourteen provide recommendations as how to screen for adolescent mental health problems (Table 12.1).

In all the countries who replied to the questionnaire ($n = 30$), it is possible for a young person to obtain emergency contraception. In about half of the countries ($n = 24$), there are multiple options where a young person can obtain emergency contraception, such as in a pharmacy, a health clinic, the emergency department of a hospital or via a primary care practitioner. 25 countries have multiple options to obtain pregnancy tests, and in most countries ($n = 24$), condoms are easily available. Only eleven countries, however, provide oral contraception free of charge, although, on the whole, adolescents can obtain such contraception easily in most countries, but in only 16 countries, it is possible for the adolescents to visit a doctor without their parents knowing. More than half of the surveyed countries ($N = 16$) have centres which provide counselling and care in sexual and reproductive health (although some centres address all ages, not specifically adolescents). In terms of primary care, however, only eight countries have specific guidelines or policies about how to address adolescent pregnancy.

Summary

Although around half of the MOCHA countries have adopted policies or guidelines that aim to secure equal access to primary care for most adolescents, including the most vulnerable, many countries of the EU and EEA lag far behind the current standards of quality health care for adolescents. The situation seems not to have improved since ten years (Ercan et al., 2009) Only a minority of countries are equipped to identify and respond to mental health emergencies and life-threatening behaviour. Access to contraception is good in most countries, but very few have developed guidelines for practitioners to help care for a pregnant adolescent. In addition, while many countries support the concept of confidential health care, only a small number provide guidelines to professionals as how to address adolescents' competence. This situation is all the more problematic as evidence suggests that the quality of primary care services has a positive effect on the health of young people (Carai, Bivol, & Chandra-Mouli, 2015; Kalamar, Bayer, & Hindin, 2016; Sanci et al., 2015). Addressing the need for specific training of health professionals is of prime importance to improve the delivery of adolescent-focused health care (see Chapter 12), and successfully addressing the complex, changing needs of adolescents (World Health Organization, 2015c).

In summary, there is a need for *all* European countries to endorse policies and strategies regarding adolescent-friendly primary care in order to improve access and quality of care for young people. The creation of specific youth clinics and addressing other important primary care services, such as public or private consultation offices and hospitals, will help to achieve these aims. No country comprehensively responds to the many facets of quality adolescent care: some have strong policies but do not secure easy access while others are in the

opposite situation. Thus, all European countries, and especially those that have a weak corpus of policies, recommendations or specific healthcare strategies (Cyprus, Hungary, Iceland, Latvia, Lithuania, Malta, Poland, Romania and Slovakia) can begin improvement in different ways.

- Physicians, especially those involved in scientific organisations or in public health activities should advocate for adolescent health, sensitising colleagues and policy-makers to the importance of this cohort. Adoption of good life-styles during this period of life will profoundly affect their health for the rest of their life (see Chapter 2).
- Addressing health-compromising behaviour, supporting healthy habits is the responsibility of adolescents' primary care providers (Patton et al., 2014, 2016; World Health Organization, 2017). European countries must develop policies and strategies which improve access to adolescents facing situations of vulnerability; particularly in the area of mental health and sexual and reproductive health. Schools, ambulatory settings and hospitals should offer easily identified, low-threshold comprehensive health care and a culturally appropriate approach, given the number of migrant adolescents being hosted in most European countries (see Chapter 11).
- Also, services to adolescents, even if they follow the evidence-based standards, will not be effective if young people themselves are inadequately informed or able to access them. It is the task of both the education and the healthcare systems to assist young people to understand their rights and responsibility for their health, and how and where to access to adequate care.
- One of the best ways to improve the quality of care delivered to adolescents is to improve the training of healthcare providers (Michaud et al., 2017). This is addressed in Chapter 13.

References

- Ambresin, A. E., Bennett, K., Patton, G. C., Sanci, L. A., & Sawyer, S. M. (2013). Assessment of youth-friendly health care: A systematic review of indicators drawn from young people's perspectives. *Journal of Adolescent Health, 52*, 670–681. doi:10.1016/j.jadohealth.2012.12.014
- Baltag, V., & Mathison, A. (2010). *Youth-friendly health policies and services in the European region: Sharing experiences* (p. 267). Copenhagen: World Health Organization.
- Bell, D. L., Breland, D. J., & Ott, M. A. (2013). Adolescent and young adult male health: A review. *Pediatrics, 132*(3), 535–546.
- Blair, M., Rigby, M., & Alexander, D. (2017). *Final report on current models of primary care for children*. Retrieved from www.childhealthservicemodels.eu/wp-content/uploads/2017/07/MOCHA-WP1-Deliverable-WP1-D6-Feb-2017-1.pdf
- Carai, S., Bivol, S., & Chandra-Mouli, V. (2015). Assessing youth-friendly-health-services and supporting planning in the Republic of Moldova. *Reproductive Health, 12*, 98.
- Committee on Adolescence. (2016). Achieving quality health services for adolescents. *Pediatrics, 138*(2).

- Ercan, O., Alikasifoglu, M., Erginoz, E., Janda, J., Kabicek, P., Rubino, A., ... Vural, M. (2009). Demography of adolescent health care delivery and training in Europe. *European Journal of Pediatrics*, 168(4), 417–426.
- Jansen, D. E. M. C., Visser, A., Vervoort, J. P. M., van der Pol, S., Kocken, P., Reijneveld, S. A., & Michaud, P. (2018). *School and adolescent health services in 30 European countries: A description of structure and functioning, and of health outcomes and costs*. Retrieved from http://www.childhealthservicemodels.eu/wp-content/uploads/Deliverable-173.1_Final-report-on-the-description-of-the-various-models-of-school-health-services-and-adolescent-health-services.pdf
- Kalamar, A. M., Bayer, A. M., & Hindin, M. J. (2016). Interventions to prevent sexually transmitted infections, including HIV, among young people in low- and middle-income countries: A systematic review of the published and gray literature. *Journal of Adolescent Health*, 59(3 Suppl), S22–31.
- Kokotailo, P. K., Baltag, V., & Sawyer, S. M. (2018). Educating and training the future adolescent health workforce. *Journal of Adolescent Health*, 62(5), 511–524.
- Michaud, P. A., Berg-Kelly, K., Macfarlane, A., & Benaroyo, L. (2010). Ethics and adolescent care: An international perspective. *Current Opinion in Pediatrics*, 22(4), 418–422.
- Michaud, P. A., Blum, R. W., Benaroyo, L., Zermatten, J., & Baltag, V. (2015). Assessing an adolescent's capacity for autonomous decision-making in clinical care. *Journal of Adolescent Health*, 57, 361–366. doi:10.1016/j.jadohealth.2015.06.012
- Michaud, P. A., Schrier, L., Ross-Russel, R., van der Heijden, L., Dossche, L., Copley, S., ... Ambresin, A. E. (2017). Paediatric departments need to improve residents' training in adolescent medicine and health: A position paper of the European Academy of Paediatrics. *European Journal of Pediatrics*, 177(4), 479–487. doi:10.1007/s00431-017-3061-2
- Michaud, P., & Baltag, V. (2015). *Core competencies in adolescent health and development for primary care providers* (p. 49). Geneva: World Health Organization.
- Michaud, P., Blum, R., & Ferron, C. (1998). "Bet you I will!" Risk or experimental behavior during adolescence? *Archives of Pediatrics & Adolescent Medicine*, 152, 224–226.
- Michaud, P., Weber, M., Namazova-Baranova, L., & Ambresin, A. (2018). Improving the quality of care delivered to adolescents in Europe: A time to invest. *Archives of Disease in Childhood*, 104(3), 214–216.
- Nair, M., Baltag, V., Bose, K., Boschi-Pinto, C., Lambrechts, T., & Mathai, M. (2015). Improving the quality of health care services for adolescents, globally: A standards-driven approach. *Journal of Adolescent Health*, 57(3), 288–298.
- National Research Council and Institute of Medicine. (2009). *Adolescent health services*. Washington, DC: National Academic Press.
- Patton, G. C., Coffey, C., Cappa, C., Currie, D., Riley, L., Gore, F., ... Ferguson, J. (2012). Health of the world's adolescents: A synthesis of internationally comparable data. *Lancet*, 379(9826), 1665–1675.
- Patton, G. C., Ross, D. A., Santelli, J. S., Sawyer, S. M., Viner, R. M., & Kleinert, S. (2014). Next steps for adolescent health: A Lancet commission. *Lancet*, 383(9915), 385–386.
- Patton, G. C., Sawyer, S. M., Santelli, J. S., Ross, D. A., Afifi, R., Allen, N. B., ... Viner, R. M. (2016). Our future: A Lancet commission on adolescent health and wellbeing. *Lancet*, 387(10036), 2423–2478.

- Potrebny, T., Wium, N., & Lundegard, M. M. (2017). Temporal trends in adolescents' self-reported psychosomatic health complaints from 1980-2016: A systematic review and meta-analysis. *PLoS One*, *12*(11), e0188374.
- Sanci, L., Chondros, P., Sawyer, S., Pirkis, J., Ozer, E., Hegarty, K., ... Patton, G. (2015). Responding to young people's health risks in primary care: A cluster randomised trial of training clinicians in screening and motivational interviewing. *PLoS One*, *10*(9), e0137581.
- Sawyer, S. M., Ambresin, A. E., Bennett, K. E., & Patton, G. C. (2014). A measurement framework for quality health care for adolescents in hospital. *Journal of Adolescent Health*, *55*(4), 484–490.
- Steinberg, L., Icenogle, G., Shulman, E. P., Breiner, K., Chein, J., Bacchini, D., ... Takash, H. M. (2017). Around the world, adolescence is a time of heightened sensation seeking and immature self-regulation. *Developmental Science*, *21*(2), e12532.
- United Nation General Assembly. (1990). *United nation convention on the rights of the child*. New York, NY: UN General Assembly.
- United Nations. (1989). *Convention on the rights of the child*, UNICEF.
- World Health Organization. (2014). *Health for the world's adolescents: A second chance in the second decade*. Geneva: World Health Organization. Retrieved from https://www.who.int/maternal_child_adolescent/documents/second-decade/en/
- World Health Organization. (2014a). *European framework for quality standards in school health services and competences for school health professionals* (Vol. 1, p. 11). Copenhagen: World Health Organization Regional Office for Europe.
- World Health Organization. (2014b). *Health for the world's adolescents: A second chance in the second decade*. Geneva: World Health Organization.
- World Health Organization. (2015a). *Children's rights in primary health care. Assessment and improvement Tool for Children and Adolescents aged 12-18* (Vol. 3, p. 50). Copenhagen: World Health Organization.
- World Health Organization. (2015b). *Core competencies in adolescent health and development for primary care providers: Including a tool to assess the adolescent health and development component in pre-service education of health-care providers* (p. 58). Geneva: World Health Organization.
- World Health Organization. (2015c). *Global standards for quality health-care services for adolescents: A guide to implement a standards-driven approach to improve the quality of health care services for adolescents*. Volume 1: Standards and criteria. Geneva: World Health Organization.
- World Health Organization. (2016). *Improving the quality of care for reproductive, maternal, neonatal, child and adolescent health in the WHO European region* (p. 50). Copenhagen: World Health Organization.
- World Health Organization. (2017). *Global accelerated action for the health of adolescents (AA-HA!): Guidance to support country implementation*. Geneva: World Health Organization.