

Seasonal variation and its impacts in rice-growing regions of the Mekong Delta

Rice-growing regions of the Mekong Delta

Van Hong Nguyen

*Sub-institute of Hydrometeorology and Climate Change (SIHYMECC),
HCM City, Vietnam, and*

Hoang Phan Hai Yen

School of Social Sciences Education, Vinh University, Nghe An, Vietnam

483

Received 22 May 2020
Revised 15 July 2020
27 December 2020
21 April 2021
26 May 2021
30 May 2021
31 May 2021
Accepted 9 June 2021

Abstract

Purpose – In recent years, agricultural activities in the Mekong Delta have frequently faced the potential risks of drought, saline intrusion and unusually heavy rainfall because of climate change, leading to a decline in crop yield. Therefore, this study aimed to establish rice planting seasons in An Giang, an upper-located province in the Mekong Delta.

Design/methodology/approach – The impacts of seasonal variation on the key rice seasons were simulated using the Food and Agriculture Organization-crop model for the OM6976 rice variety grown in the study area. For the simulation, the model combined crop, soil, weather and crop management data.

Findings – The results show that seasonal variation because of changes in weather factors leads to alternation in crop yields across the study area. Specifically, the spring and summer rice planting seasons are advanced by one to two weeks compared with the baseline, and crop yield increased by 5.9% and 4.2%, respectively. Additionally, planting for the autumn–winter rice season on 3 August increased crop yield by up to 8.1%.

Originality/value – In general, rice planting seasons that account for weather factor changes effectively reduce production costs and optimise production.

Keywords Crop, Seasonal variation, Weather factors, Cultivation, Mekong Delta

Paper type Research paper

1. Introduction

In the last two decades, under the impact of climate variability (ICV), rainfall has changed significantly in various regions of the world (APN, 2010; Kontgis *et al.*, 2019; Furuya and Koyama, 2005), leading to a lack of water for irrigation activities in the dry season, flooding in the rainy season (Brauman *et al.*, 2013) and extreme rainfall events causing losses after

© Van Hong Nguyen and Hoang Phan Hai Yen. Published by Emerald Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

Author contributions: H.P.H. Yen and N.V. Hong discussed the original idea of the draft. H.P.H. Yen analysed and designed the input data. N.V. Hong established the input data and performed the model simulation. H.P.H. Yen wrote and edited the manuscript. N.V. Hong analysed the output data. Both authors reviewed and submitted the final version of the manuscript.

Conflict of interest: The authors declare that there are no conflicts of interest.



International Journal of Climate
Change Strategies and
Management
Vol. 13 No. 4/5, 2021
pp. 483-491
Emerald Publishing Limited
1756-8692
DOI 10.1108/IJCCSM-05-2020-0048

harvest (Deryng *et al.*, 2014; Zhu and Troy, 2018). The RCSA (2016) reported that climate variability would impact many Asian regions, including the Mekong Delta, in the 2020s. Climate variability is one of the main causes of drought, leading to water scarcity and unseasonal rainfall, which are the major factors limiting crop yields. According to Croitoru, rainfall is an important factor that can be directly affected by climate variability. In this context, compliance with agronomic practices such as fertiliser application, suitable plant densities, adoption of appropriate rice varieties (Bai and Xiao, 2019) and timeous planting (Dang, 2021; Trang, 2016) has been considered optimal solutions. Agronomic practices help minimise the negative ICV (Dang, 2021; Lobell, 2014; Giorgi *et al.*, 2016) and contribute to improving crop yields (Leng and Huang, 2017). According to Sacks, selecting adaptable cultivars and planting are two of the top criteria that help to increase crop yields in the context of climate variability (CCV).

In recognition of the importance of seasonal variation in rice-growing regions in the CCV, a study on the changes in the planting seasons as an adaptation solution for rice production in Kurunegala was conducted by Dharmarathna *et al.* (2014). They concluded that the crop yield in the dry season would increase compared with the current crop calendar if the planting seasons were delayed by four weeks under the regional growth scenario (A2) and the increasing population scenario (B2). In the Ca Mau Peninsula of Vietnam, Deb *et al.* (2015) reported that a shift in planting seasons was beneficial for enhancing rice yields under the ICV. In 2016, Shrestha *et al.* (2016) conducted a study on adaptation strategies for rice-growing regions in Central Vietnam under the ICV and reported that a delay in planting resulted in a projected rice yield increase of up to 20%. In 2017, Li *et al.* (2017) reported that advancing planting by three weeks compared with the baseline would slightly increase rice yields in Cambodia by 0%–4.7%.

Vietnam has faced a high risk of crop failure because of the ICV in recent years (MNRE, 2016). Farmers usually rely on irrigation water from the Mekong River and local rainfall (Mainuddin *et al.*, 2013). These crops face regular potential risks and must be replanted if heavy rainfall events occur after seeds have been planted or during the flowering and harvesting stages. Salim (2009) recommended that rice planting seasons in Vietnam should be altered to ensure optimal grain yield under the ICV. The harvesting stage of the summer crop season often coincides with peak-intensity rainfall varying from 10 to 50 mm/day, leading to a significant reduction in the harvested yield owing to rice paddy collapse following heavy rainfall events.

In the context of seasonal variation, the ICV has negatively affected crop seasons, so planting seasons need to be re-delineated to ensure that the different rice crop growth stages receive appropriate rainwater irrigation, contributing to minimising the adverse effects of weather factors as well as improving grain yield in the study area. Therefore, by focusing on rice paddies across An Giang Province as a typical case study, this study aimed to highlight the role of weather factors in crop productivity through a combination of statistical analysis and crop model simulation.


2. Materials and methods

2.1 Study area

An Giang, an upper-located province in the Mekong Delta, lies between 10°12' N latitude and 10°57' N and 104°46' E and 105°35' E longitude (Figure 1). It has a total agricultural land area of approximately 300,000 ha, with elevations ranging from 0.5–2.5 m a.m.s.l (Dang, 2021; Vu *et al.*, 2018). An Giang is an agricultural province known to have the highest production in Vietnam, with average rice productivity reaching 6.74 ton per ha (Trang, 2016). Agriculture is considered a key sector, with rice production based on soil

Figure 1.

Map of the study area
with rainfall
observation stations
for conducting the
model calibration and
validation marked the
green circles



Source: Figure is created by the authors

fertility and freshwater from the upper Mekong River and local rainfall (Vo and Huynh, 2014).

In An Giang Province, farmers often have three rice crop seasons per year, including spring, summer and autumn–winter planting seasons. The flowering and harvesting stages can vary from season to season and year to year based on weather conditions, as well as the decisions of professional management agencies. The East Asian monsoon circulation is dominant in the study area and is divided into two main seasons: the southwest and northeast monsoon. The southwest monsoon is characterised by heat and humidity with abundant rainfall, whereas the northeast monsoon is dry and hot, producing very little rainfall. Average temperatures range between 26.3°C and 28.5°C, with average annual precipitation varying from 1,268.5 mm at the Chau Doc station to 1,549.6 mm at the Long Xuyen station, of which approximately 90% falls in the wet season (Table 1).

Rainfall is highly concentrated in the western and southern districts, with up to 1,430 mm per year, and it decreases gradually in the eastern and northern districts, down to 1,350 mm per year. In recent years, a significant reduction in rainfall has been recorded in the northern districts, which could negatively impact agricultural production in the study area (Vo and Huynh, 2014). Therefore, the area faces the ICV, leading to drought, flooding and unseasonal rainfall. For example, in 2016, the worst drought event in 90 years was recorded in Vietnam;

Table 1.
Average annual
rainfall (AAR) and
standard deviation
(SD) at weather
stations

No.	Station name	AAR (mm)	SD (mm)	Latitude (N)	Longitude (E)
1	Chau Doc	1,285.6	76.6	10°33'24	105°08'09
2	Xuan To	1,467.9	84.7	10°35'32	104°56'44
3	Tri Ton	1,421.4	85.0	10°23'50	104°59'08
4	Long Xuyen	1,399.8	97.6	10°23'08	105°26'04
5	Tan Chau	1,300.1	88.7	10°50'32	105°11'04
6	Phu Tan	1,378.7	86.2	10°40'00	105°17'22
7	Cho Moi	1,409.6	86.5	10°28'57	105°28'37

the lower Mekong River receded to its lowest level since 1926, and water shortages hampered agricultural productivity in regions of Vietnam (FAO, 2016). Furthermore, the collapse of the Xe-PianXe-Namnoy dam in Laos in 2018 because of heavy rainfall led to rapidly increasing floodwater levels which impeded the ongoing harvest of the summer crop (Mainuddin *et al.*, 2013). The low-lying geography of the study area makes it vulnerable to sea-level rise, salinity intrusion and flooding because of heavy rainfall events (RCSA, 2016). The area is divided into two subregions (Figure 1) based on topographical conditions and irrigation water sources. Specifically, subregion I is characterised by a lower topography from north to south and east to west, with the rice paddies receiving irrigation water from the Hau River and local rainfall. However, the rice fields in the coastal districts sometimes regularly face saline intrusion during the dry season and flooding during the wet season. Subregion II is characterised by a lower topography from north to south, and the rice fields receive irrigation water from the Tien and Hau Rivers and local rainfall. Subregion II has not faced saline intrusion and is rarely flooded during the wet season.


2.2 Model description

The Food and Agriculture Organization (FAO)-crop model was designed by the FAO to evaluate the irrigation water requirement, biomass, component and crop yields, fertiliser, nutrient management and planting under current and future climatic conditions (Lee and Dang, 2019; Steduto *et al.*, 2009). Its performance has been demonstrated by study outcomes that have been widely adopted for a wide range of applications at different spatial and temporal scales, for example, simulating a maize crop in a semi-arid region in India; predicting the spatiotemporal shifts of rice phenology in China from 1981–2010 (Bai and Xiao, 2019); estimating rice productivity in the Ca Mau Peninsula, Vietnam (Deb *et al.*, 2015); and simulating canopy cover, biomass, yield and water requirement in Southern Taiwan (Greaves and Wang, 2016). Details of the crop model can be found in FAO, 2015.

2.3 Data collection

To conduct this study, weather data from 2000 to 2018 were obtained from Phu Tan and Long Xuyen meteorological stations, whereas evaporation data were calculated based on the Penman-Monteith formula. Specifically, rainfall data from 2000 to 2010 [Figure 2(a)] were

Figure 2.
Precipitation data for
a) model calibration
using 2000–2010 data
and b) model
validation using
2011–2018 data



Source: Figure is created by the authors

used for model calibration, whereas rainfall data from 2011 to 2018 [Figure 2(b)] were applied for model validation.

To simulate crop yield, detailed data on the soil characteristics were collected from a field survey at Phu Tan and Long Xuyen stations and were analysed for application as input data. The analysis indicated that the soil components were sandy loam to loamy sand and alkaline to slightly alkaline ($\text{pH} = 4.6\text{--}5.8$). The soil contained a Ca^{2+} content of $7.5\text{--}13.0 \text{ mg kg}^{-1}$ and a Mg^{2+} content of $2.69\text{--}4.73 \text{ mg kg}^{-1}$ in the upper surface layer (0–120 cm). Based on the indicative values for soil water content by the FAO (2015), the moisture content at saturation (SAT) was 47%, field capacity was 32%, permanent wilting point was approximately 20%, saturated hydraulic conductivity was 120 mm m^{-1} and total available soil water was 225 mm per day. The crop management data for rice cultivation were collected from the Department of Agriculture and Rural Development. Detailed information on planting, harvesting times, density at planting, fertiliser application and irrigation amount are presented in Table 2. OM6976 is a rice cultivar with high productivity and good pest resistance, with a planting cycle varying from 95 to 105 days and is planted widely in the area (Table 2).

The paddies are commonly planted in the first week of December and harvested in the third week of March for the spring rice season, planted in the second week of April and harvested in the fourth week of July for the summer rice season and planted in the second week of August and harvested in the fourth week of November for the autumn–winter rice season. Fertiliser was applied at the recommended rates to obtain an optimal crop yield for all crop planting seasons varying from $80\text{--}100 \text{ kg ha}^{-1}$ for Urea, $60\text{--}70 \text{ kg ha}^{-1}$ for N and 60 kg ha^{-1} for K. The optimal fertiliser schedules were determined for the first time as 8–12 days after planting (DAP), the second time at 18–22 DAP, the third time at 27–32 DAP and the final time at 40–43 DAP (Table 2).

2.4 Model performance assessment

The model performance was assessed using the agreement of regression equations based on the observation data and the calculated model. Error statistics such as the coefficient of determination (R^2), index of agreement (d) and root mean square error (RMSE) were used in this study (Greaves and Wang, 2016; Lee and Dang, 2019).

3. Results and discussion

3.1 Model calibration and validation

The crop model was calibrated using the rice yield of the three rice crops during 2000–2010 period. Results indicated a strong linear correlation between the model and the observed yield corresponding to the spring, summer and autumn–winter planting seasons [Figure 3 (a), 3(b), 3(c)] in subregion I, with d , R^2 and RMSE varying from 0.84–0.93, 0.89–0.94 and

Crop	Crop calendar		Variety	Density (kg ha^{-1})	Fertiliser schedules and applications rate						
	Plant	Harvest			I	II	III	IV	Urea	N	K
Spring	05 December	15 March	OM6976	180	10	20	28	42	80	70	60
Summer	08 April	24 July	OM6976	200	8	16	27	36	90	70	60
Autumn–winter	10 August	22 November	OM6976	180	9	17	26	35	100	60	60

Notes: I, II, III and IV are days after planting (DAP)

Table 2.
Crop calendar,
fertiliser schedules
and application rate
for rice varieties

Figure 3.
Results of model
calibration based on
the simulated and
observed yield of a)
spring, b) summer
and c) autumn–
winter planting
seasons in subregion
I; and d) spring, e)
summer and f)
autumn–winter
planting seasons in
subregion II

0.17–0.38, respectively; the corresponding values for subregion II were 0.79–0.88, 0.86–0.91 and 0.21–0.45, respectively (Table 3).

Similarly, model validation was also performed for the three rice crop planting seasons using crop yield from the 2011–2018 period. The accuracy of the applied model is shown in Table 3. Results showed a high correlation between the model and observed yield with d , R^2 , and RMSE varying from 0.87–0.92, 0.91–0.95 and 0.19–0.27, respectively, in subregion I; the corresponding values for subregion II were 0.83–0.90, 0.87–0.91 and 0.24–0.37, respectively Figure 4.

3.2 Impact of seasonal variation on crop yield

For years, rice has been cultivated in An Giang Province without considering seasonal variations. Cultivating rice under inappropriate weather conditions, such as high daily rainfall during harvest, can reduce rice production. According to Kunimistu and Kudo (2015), heavy rainfall events that occur continuously during the flowering stage of rice can reduce the grain filling process.

The results indicated that the crop yield of rice planting seasons in the An Giang Province increased notably if the rice received low daily rainfall during the flowering and harvesting stages. Specifically, for spring and summer planting seasons, a planting delay of one to two weeks resulted in low daily rainfall during the flowering and harvesting stages. Crop yield of the spring and summer planting seasons increased by approximately 5.9% and 5.1% for subregions I and II, respectively (Table 4).




Table 3.
Evaluation results of
the performance of
the applied model

Region	Crop	Calibration			Validation		
		d	RMSE	R^2	d	RMSE	R^2
Subregion I	Spring	0.93	0.17	0.94	0.92	0.19	0.95
	Summer	0.84	0.38	0.89	0.87	0.33	0.91
	Autumn–winter	0.88	0.25	0.91	0.89	0.27	0.93
Subregion II	Spring	0.88	0.21	0.91	0.90	0.24	0.91
	Summer	0.79	0.45	0.86	0.83	0.37	0.87
	Autumn–winter	0.82	0.31	0.89	0.85	0.29	0.90




Figure 4. Results of the model validation based on the simulated and observed yield of a) spring, b) summer and c) autumn–winter planting seasons in subregion I; and d) spring, e) summer and f) autumn–winter planting seasons in subregion II

Source: Figure is created by the authors

		Percentage yield change corresponds to the different planting seasons										
		Earlier (week)					Later (week)					
Region	Crop	−5	−4	−3	−2	−1	Current (ton ha ^{−1})	1	2	3	4	5
I	Spring	−5.9	−4.4	−2.9	−2.9	1.7	6.8	4.3	5.9	4.4	2.9	−1.5
	Summer	−4.8	−6.3	−7.9	−4.8	−3.2	6.3	3.2	4.2	3.2	3.2	−1.6
	Autumn-winter	−1.5	1.5	3.0	6.1	3.0	6.6	2.1	−3.0	−6.1	−4.5	−2.8
II	Spring	−6.2	−4.6	−4.6	−3.1	−1.5	6.5	4.6	3.1	4.2	1.5	−1.3
	Summer	−1.7	−3.4	−5.1	−1.7	−1.7	5.9	5.1	5.1	5.1	3.4	−1.4
	Autumn-winter	−1.6	3.2	3.2	8.1	3.2	6.2	2.0	−4.8	−6.5	−4.8	−2.6

Table 4. Change in rice yield (%) for different planting times in the study area

Consequently, if planting is delayed by two weeks compared with the baseline, the autumn–winter planting season can approach the maximum crop yield. This implies that bringing forward the autumn–winter planting season by one to two weeks will help achieve an optimal crop yield. These simulated results are consistent with the actual weather conditions. When planting is brought forward by one to two weeks, rice will receive high daily rainfall during its growth phase but will receive less daily rainfall during the flowering and harvesting phases, which is favorable for high-yielding crops.

4. Conclusion

The impacts of seasonal variation due to climate change in the rice cultivating regions across the An Giang Province were estimated using the FAO-crop model.

The model was applied to simulate the crop yield of three key rice planting seasons in the An Giang Province based on combining crop, soil, weather and crop management data. The results show that the crop yield of the three rice seasons can be significantly influenced by seasonal variation because of climate change. In general, the planting calendar of the three rice seasons is no longer suitable, and farmers are exposed to high risks leading to crop failure. Therefore, it is necessary to adopt technological solutions in the near future to enhance rice production abilities and mitigate the effects of global climate variability.

References

- APN-Asia Pacific Network (2010), "Climate change in Southeast Asia and assessment on impact, vulnerability and adaptation on rice production and water resource", Project Reference Number: CRP2008-03CMY-Jintrawet.
- Bai, H. and Xiao, D. (2019), "Spatiotemporal changes of rice phenology in China during 1981–2010", *Theoretical and Applied Climatology*, Vol. 157, pp. 1483-1494.
- Brauman, K.A., Siebert, S. and Foley, J.A. (2013), "Improvements in crop water productivity increase water sustainability and food security – a global analysis", *Environmental Research Letters*, Vol. 8 No. 2, pp. 1-7.
- Dang, T.A. (2021), "Grain yield optimisation in the plain of reeds in the context of climate variability", *Brazilian Journal of Agricultural and Environmental Engineering*, Vol. 25 No. 9, pp. 591-596.
- Deb, P., Tran, D.A. and Udmale, P.D. (2015), "Assessment of the impacts of climate change and brackish irrigation water on rice productivity and evaluation of adaptation measures in Ca Mau province, Vietnam", *Theoretical and Applied Climatology*, Vol. 125 Nos 3/4, pp. 641-656.
- Deryng, D., Conway, D., Ramankutty, N., Price, J. and Warren, R. (2014), "Global crop yield response to extreme heat stress under multiple climate change futures", *Environmental Research Letters*, Vol. 9 No. 3, pp. 1-13.
- Dharmarathna, W.R.S.S., Herath, S. and Weerakoon, S.B. (2014), "Changing the planting date as a climate change adaptation strategy for rice production in Kurunegala district, Sri Lanka", *Sustainability Science*, Vol. 9 No. 1, pp. 103-111.
- FAO- Food and Agriculture Organization (2015), "AquaCrop new features and updates version 5.0", October 2015.
- FAO- Food and Agriculture Organization (2016), "El niño event in Viet Nam: Agriculture, food security and livelihood need assessment in response to drought and salt-water intrusion", Assessment Report, p. 75.
- Furuya, J. and Koyama, O. (2005), "Impacts of climatic change on world agricultural product markets: estimation of macro yield functions", *Japan Agricultural Research Quarterly: JARQ*, Vol. 39 No. 2, pp. 121-134.
- Giorgi, F., Torma, C., Coppola, E., Ban, N., Schär, C. and Somot, S. (2016), "Enhanced summer convective rainfall at alpine high elevations in response to climate warming", *Nature Geoscience*, Vol. 9 No. 8, pp. 584-589.
- Greaves, G.E. and Wang, Y.M. (2016), "Assessment of FAO AquaCrop model for simulating maize growth and productivity under deficit irrigation in a tropical environment", *Water*, Vol. 8 No. 12, pp. 2-18.
- Kontgis, C., Schneider, A., Ozdogan, M., Kucharik, C., Pham, V.D.T., Nguyen, H.D. and Schatz, Z. (2019), "Climate change impacts on rice productivity in the Mekong river Delta", *Applied Geography*, Vol. 102, pp. 71-83.
- Kunimistu, Y.J. and Kudo, R.J. (2015), "Fluctuations in rice productivity caused by long and heavy rain under climate change in Japan: evidence from panel data regression analysis", *Japan Agricultural Research Quarterly: Jarq*, Vol. 49 No. 2, pp. 159-172.

- Lee, S.K. and Dang, T.A. (2019), "Calibration and validation of the FAO-AquaCrop model for cassava in the dong Xuan cultivation area of Phu Yen province using irrigation rainfall", *Res. on Crops*, Vol. 20 No. 3, pp. 555-562.
- Leng, G. and Huang, M. (2017), "Crop yield response to climate change varies with crop spatial distribution pattern", *Scientific Reports*, Vol. 7 No. 1, pp. 1-10.
- Li, S., Wang, A. and Chun, J.A. (2017), "Impact assessment of climate change on rice productivity in the Indochinese peninsula using a regional-scale crop model", *International Journal of Climatology*, Vol. 37 No. 1, pp. 1147-1160.
- Lobell, D.B. (2014), "Climate change adaptation in crop production beware of illusions", *Global Food Security*, Vol. 3 No. 2, pp. 72-76.
- Mainuddin, M., Kirby, M. and Hoanh, C.T. (2013), "Impact of climate change on rainfed rice and options for adaptation in the lower Mekong basin", *Natural Hazards*, Vol. 66 No. 2, pp. 905-938.
- MNRE-Ministry of Natural Resources and Environment (2016), *Climate Change Scenarios and Sea Level Rise for Vietnam*, Publishers resources, environment and map of Vietnam.
- RCSA-Research Centers in Southeast Asia (2016), "The drought and salinity intrusion in the Mekong river Delta of Vietnam", Assessment Report, p. 55.
- Shrestha, S., Deb, P. and Bui, T.T.T. (2016), "Adaptation strategies for rice cultivation under climate change in Central vietnam", *Mitigation and Adaptation Strategies for Global Change*, Vol. 21 No. 1, pp. 15-37.
- Steduto, P., Hsiao, T.C., Raes, D. and Fereres, E. (2009), "AquaCrop-the FAO crop model to simulate yield response to water", *Agronomy Journal*, Vol. 101 Issue No. 3, pp. 426-437.
- Trang, T.H.L. (2016), "Effects of climate change on rice yield and rice market in Vietnam", *Journal of Agricultural and Applied Economics*, Vol. 48 No. 4, pp. 366-382.
- Vo, T.D. and Huynh, V.K. (2014), "Using a risk cost-benefit analysis for a sea dike to adapt to the sea level in the Vietnamese Mekong river Delta", *Climate*, Vol. 2 No. 2, pp. 78-102.
- Vu, D.T., Yamada, T. and Ishidaira, H. (2018), "Assessing the impact of sea level rise due to climate change on seawater intrusion in Mekong Delta, Vietnam", *Water Science and Technology*, Vol. 77 No. 6, pp. 1632-1639.
- Zhu, X. and Troy, T.J. (2018), "Agriculturally relevant climate extremes and their trends in the world's major planting regions", *Earth's Future*, Vol. 6 No. 4, pp. 656-672.

Further reading

- Sacks, W.J., Deryng, D., Foley, J.A. and Ramankutty, N. (2010), "Crop planting dates: an analysis of global patterns", *Global Ecology and Biogeography*, Vol. 19 No. 5, pp. 607-620.

Corresponding author

Hoang Phan Hai Yen can be contacted at: hoangphanhaiyen@vinhuni.edu.vn

For instructions on how to order reprints of this article, please visit our website:

www.emeraldgrouppublishing.com/licensing/reprints.htm

Or contact us for further details: permissions@emeraldinsight.com