

```

NEW FILE.
DATASET NAME DataSet1 WINDOW=FRONT.
GET DATA /TYPE=XLSX
  /FILE='I:\spss.xlsx'   /SHEET=name 'Φύλλο1'
  /CELLRANGE=full
  /READNAMES=on
  /ASSUMEDSTRWIDTH=32767.
EXECUTE.
DATASET NAME DataSet2 WINDOW=FRONT.
CROSSTABS
  /TABLES=R1D1 BY R2D1
  /FORMAT=AVALUE TABLES
  /STATISTICS=KAPPA CHISQ CORR
  /CELLS=COUNT ROW COLUMN TOTAL.

```

Crosstabs

Notes

Output Created		24-SEP-2025 14:53:41
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	112
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax		CROSSTABS /TABLES=R1D1 BY R2D1 /FORMAT=AVALUE TABLES /STATISTICS=KAPPA CHISQ CORR /CELLS=COUNT ROW COLUMN TOTAL.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,02
	Dimensions Requested	2
	Cells Available	524245

[DataSet2]

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
R1D1 * R2D1	112	100,0%	0	0,0%	112	100,0%

R1D1 * R2D1 Crosstabulation

			R2D1		Total
			,0	1,0	
R1D1	,0	Count	53	5	58
		% within R1D1	91,4%	8,6%	100,0%
		% within R2D1	84,1%	10,2%	51,8%
		% of Total	47,3%	4,5%	51,8%
	1,0	Count	10	44	54
		% within R1D1	18,5%	81,5%	100,0%
		% within R2D1	15,9%	89,8%	48,2%
		% of Total	8,9%	39,3%	48,2%
Total		Count	63	49	112
		% within R1D1	56,3%	43,8%	100,0%
		% within R2D1	100,0%	100,0%	100,0%
		% of Total	56,3%	43,8%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	60,324 ^a	1	,000	,000	,000
Continuity Correction ^b	57,400	1	,000		
Likelihood Ratio	67,694	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	59,786	1	,000		
N of Valid Cases	112				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 23,63.

b. Computed only for a 2x2 table

Symmetric Measures

		Value	Asymptotic Standardized Error ^a	Approximate T ^b	Approximate Significance
Interval by Interval	Pearson's R	,734	,063	11,332	,000 ^c
Ordinal by Ordinal	Spearman Correlation	,734	,063	11,332	,000 ^c
Measure of Agreement	Kappa	,731	,064	7,767	,000
N of Valid Cases		112			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

CROSSTABS

/TABLES=R1D2 BY R2D2

/FORMAT=AVALUE TABLES

/STATISTICS=KAPPA CHISQ CORR

/CELLS=COUNT ROW COLUMN TOTAL.

Crosstabs

Notes

Output Created		24-SEP-2025 14:53:41
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	112
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax		CROSSTABS /TABLES=R1D2 BY R2D2 /FORMAT=AVALUE TABLES /STATISTICS=KAPPA CHISQ CORR /CELLS=COUNT ROW COLUMN TOTAL.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,01
	Dimensions Requested	2
	Cells Available	524245

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
R1D2 * R2D2	112	100,0%	0	0,0%	112	100,0%

R1D2 * R2D2 Crosstabulation

			R2D2		Total
			,0	1,0	
R1D2	,0	Count	85	4	89
		% within R1D2	95,5%	4,5%	100,0%
		% within R2D2	97,7%	16,0%	79,5%
		% of Total	75,9%	3,6%	79,5%
	1,0	Count	2	21	23
		% within R1D2	8,7%	91,3%	100,0%
		% within R2D2	2,3%	84,0%	20,5%
		% of Total	1,8%	18,8%	20,5%
Total	Count		87	25	112
	% within R1D2		77,7%	22,3%	100,0%
	% within R2D2		100,0%	100,0%	100,0%
	% of Total		77,7%	22,3%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	79,436 ^a	1	,000	,000	,000
Continuity Correction ^b	74,508	1	,000		
Likelihood Ratio	72,706	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	78,726	1	,000		
N of Valid Cases	112				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 5,13.

b. Computed only for a 2x2 table

Symmetric Measures

		Value	Asymptotic Standardized Error ^a	Approximate T ^b	Approximate Significance
Interval by Interval	Pearson's R	,842	,062	16,381	,000 ^c
Ordinal by Ordinal	Spearman Correlation	,842	,062	16,381	,000 ^c
Measure of Agreement	Kappa	,841	,063	8,913	,000
N of Valid Cases		112			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

CROSSTABS

/TABLES=R1D3 BY R2D3

/FORMAT=AVALUE TABLES

/STATISTICS=KAPPA CHISQ CORR

/CELLS=COUNT ROW COLUMN TOTAL.

Crosstabs

Notes

Output Created		24-SEP-2025 14:53:41
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	112
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax		CROSSTABS /TABLES=R1D3 BY R2D3 /FORMAT=AVALUE TABLES /STATISTICS=KAPPA CHISQ CORR /CELLS=COUNT ROW COLUMN TOTAL.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,02
	Dimensions Requested	2
	Cells Available	524245

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
R1D3 * R2D3	112	100,0%	0	0,0%	112	100,0%

R1D3 * R2D3 Crosstabulation

			R2D3		Total
			,0	1,0	
R1D3	,0	Count	39	0	39
		% within R1D3	100,0%	0,0%	100,0%
		% within R2D3	97,5%	0,0%	34,8%
		% of Total	34,8%	0,0%	34,8%
	1,0	Count	1	72	73
		% within R1D3	1,4%	98,6%	100,0%
		% within R2D3	2,5%	100,0%	65,2%
		% of Total	0,9%	64,3%	65,2%
Total		Count	40	72	112
		% within R1D3	35,7%	64,3%	100,0%
		% within R2D3	100,0%	100,0%	100,0%
		% of Total	35,7%	64,3%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	107,704 ^a	1	,000	,000	,000
Continuity Correction ^b	103,451	1	,000		
Likelihood Ratio	135,426	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	106,742	1	,000		
N of Valid Cases	112				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 13,93.

b. Computed only for a 2x2 table

Symmetric Measures

		Value	Asymptotic Standardized Error ^a	Approximate T ^b	Approximate Significance
Interval by Interval	Pearson's R	,981	,019	52,515	,000 ^c
Ordinal by Ordinal	Spearman Correlation	,981	,019	52,515	,000 ^c
Measure of Agreement	Kappa	,980	,019	10,378	,000
N of Valid Cases		112			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

CROSSTABS

/TABLES=R1D4 BY R2D4

/FORMAT=AVALUE TABLES

/STATISTICS=KAPPA CHISQ CORR

/CELLS=COUNT ROW COLUMN TOTAL.

Crosstabs

Notes

Output Created		24-SEP-2025 14:53:41
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	112
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax		CROSSTABS /TABLES=R1D4 BY R2D4 /FORMAT=AVALUE TABLES /STATISTICS=KAPPA CHISQ CORR /CELLS=COUNT ROW COLUMN TOTAL.
Resources	Processor Time	00:00:00,00
	Elapsed Time	00:00:00,01
	Dimensions Requested	2
	Cells Available	524245

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
R1D4 * R2D4	112	100,0%	0	0,0%	112	100,0%

R1D4 * R2D4 Crosstabulation

			R2D4		Total
			,0	1,0	
R1D4	,0	Count	95	0	95
		% within R1D4	100,0%	0,0%	100,0%
		% within R2D4	97,9%	0,0%	84,8%
		% of Total	84,8%	0,0%	84,8%
	1,0	Count	2	15	17
		% within R1D4	11,8%	88,2%	100,0%
		% within R2D4	2,1%	100,0%	15,2%
		% of Total	1,8%	13,4%	15,2%
Total	Count		97	15	112
	% within R1D4		86,6%	13,4%	100,0%
	% within R2D4		100,0%	100,0%	100,0%
	% of Total		86,6%	13,4%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	96,786 ^a	1	,000	,000	,000
Continuity Correction ^b	89,328	1	,000		
Likelihood Ratio	75,893	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	95,922	1	,000		
N of Valid Cases	112				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,28.

b. Computed only for a 2x2 table

Symmetric Measures

		Value	Asymptotic Standardized Error ^a	Approximate T ^b	Approximate Significance
Interval by Interval	Pearson's R	,930	,048	26,453	,000 ^c
Ordinal by Ordinal	Spearman Correlation	,930	,048	26,453	,000 ^c
Measure of Agreement	Kappa	,927	,051	9,838	,000
N of Valid Cases		112			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

CROSSTABS

/TABLES=R1D5 BY R2D5

/FORMAT=AVALUE TABLES

/STATISTICS=KAPPA CHISQ CORR

/CELLS=COUNT ROW COLUMN TOTAL.

Crosstabs

Notes

Output Created		24-SEP-2025 14:53:41
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	112
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax		CROSSTABS /TABLES=R1D5 BY R2D5 /FORMAT=AVALUE TABLES /STATISTICS=KAPPA CHISQ CORR /CELLS=COUNT ROW COLUMN TOTAL.
Resources	Processor Time	00:00:00,00
	Elapsed Time	00:00:00,00
	Dimensions Requested	2
	Cells Available	524245

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
R1D5 * R2D5	112	100,0%	0	0,0%	112	100,0%

R1D5 * R2D5 Crosstabulation

			R2D5		Total
			,0	1,0	
R1D5	,0	Count	6	1	7
		% within R1D5	85,7%	14,3%	100,0%
		% within R2D5	85,7%	1,0%	6,3%
		% of Total	5,4%	0,9%	6,3%
	1,0	Count	1	104	105
		% within R1D5	1,0%	99,0%	100,0%
		% within R2D5	14,3%	99,0%	93,8%
		% of Total	0,9%	92,9%	93,8%
Total		Count	7	105	112
		% within R1D5	6,3%	93,8%	100,0%
		% within R2D5	100,0%	100,0%	100,0%
		% of Total	6,3%	93,8%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	80,467 ^a	1	,000	,000	,000
Continuity Correction ^b	66,651	1	,000		
Likelihood Ratio	35,329	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	79,749	1	,000		
N of Valid Cases	112				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is ,44.

b. Computed only for a 2x2 table

Symmetric Measures

		Value	Asymptotic Standardized Error ^a	Approximate T ^b	Approximate Significance
Interval by Interval	Pearson's R	,848	,106	16,754	,000 ^c
Ordinal by Ordinal	Spearman Correlation	,848	,106	16,754	,000 ^c
Measure of Agreement	Kappa	,848	,106	8,970	,000
N of Valid Cases		112			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

CROSSTABS

/TABLES=R1D6 BY R2D6

/FORMAT=AVALUE TABLES

/STATISTICS=KAPPA CHISQ CORR

/CELLS=COUNT ROW COLUMN TOTAL.

Crosstabs

Notes

Output Created		24-SEP-2025 14:53:41
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	112
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax		CROSSTABS /TABLES=R1D6 BY R2D6 /FORMAT=AVALUE TABLES /STATISTICS=KAPPA CHISQ CORR /CELLS=COUNT ROW COLUMN TOTAL.
Resources	Processor Time	00:00:00,00
	Elapsed Time	00:00:00,00
	Dimensions Requested	2
	Cells Available	524245

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
R1D6 * R2D6	112	100,0%	0	0,0%	112	100,0%

R1D6 * R2D6 Crosstabulation

			R2D6		Total
			,0	1,0	
R1D6	,0	Count	89	0	89
		% within R1D6	100,0%	0,0%	100,0%
		% within R2D6	96,7%	0,0%	79,5%
		% of Total	79,5%	0,0%	79,5%
	1,0	Count	3	20	23
		% within R1D6	13,0%	87,0%	100,0%
		% within R2D6	3,3%	100,0%	20,5%
		% of Total	2,7%	17,9%	20,5%
Total		Count	92	20	112
		% within R1D6	82,1%	17,9%	100,0%
		% within R2D6	100,0%	100,0%	100,0%
		% of Total	82,1%	17,9%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	94,216 ^a	1	,000	,000	,000
Continuity Correction ^b	88,381	1	,000		
Likelihood Ratio	87,294	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	93,374	1	,000		
N of Valid Cases	112				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,11.

b. Computed only for a 2x2 table

Symmetric Measures

		Value	Asymptotic Standardized Error ^a	Approximate T ^b	Approximate Significance
Interval by Interval	Pearson's R	,917	,045	24,140	,000 ^c
Ordinal by Ordinal	Spearman Correlation	,917	,045	24,140	,000 ^c
Measure of Agreement	Kappa	,914	,049	9,706	,000
N of Valid Cases		112			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

CROSSTABS

/TABLES=R1D7 BY R2D7

/FORMAT=AVALUE TABLES

/STATISTICS=KAPPA CHISQ CORR

/CELLS=COUNT ROW COLUMN TOTAL.

Crosstabs

Notes

Output Created	24-SEP-2025 14:53:41	
Comments		
Input	Active Dataset	DataSet2
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	112
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax	CROSSTABS /TABLES=R1D7 BY R2D7 /FORMAT=AVALUE TABLES /STATISTICS=KAPPA CHISQ CORR /CELLS=COUNT ROW COLUMN TOTAL.	
Resources	Processor Time	00:00:00,00
	Elapsed Time	00:00:00,00
	Dimensions Requested	2
	Cells Available	524245

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
R1D7 * R2D7	112	100,0%	0	0,0%	112	100,0%

R1D7 * R2D7 Crosstabulation

			R2D7		Total
			,0	1,0	
R1D7	,0	Count	99	3	102
		% within R1D7	97,1%	2,9%	100,0%
		% within R2D7	99,0%	25,0%	91,1%
		% of Total	88,4%	2,7%	91,1%
	1,0	Count	1	9	10
		% within R1D7	10,0%	90,0%	100,0%
		% within R2D7	1,0%	75,0%	8,9%
		% of Total	0,9%	8,0%	8,9%
Total	Count		100	12	112
	% within R1D7		89,3%	10,7%	100,0%
	% within R2D7		100,0%	100,0%	100,0%
	% of Total		89,3%	10,7%	100,0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	72,154 ^a	1	,000	,000	,000
Continuity Correction ^b	63,341	1	,000		
Likelihood Ratio	42,701	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	71,510	1	,000		
N of Valid Cases	112				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 1,07.

b. Computed only for a 2x2 table

Symmetric Measures

		Value	Asymptotic Standardized Error ^a	Approximate T ^b	Approximate Significance
Interval by Interval	Pearson's R	,803	,094	14,114	,000 ^c
Ordinal by Ordinal	Spearman Correlation	,803	,094	14,114	,000 ^c
Measure of Agreement	Kappa	,799	,097	8,494	,000
N of Valid Cases		112			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.